

Chemical Functional Physical and Sensory Properties of Flour and Idli Produced from Blends of Rice (*Oryza Sativa*), African Yam Bean (*Sphenostylis sternocarpa*) and Pigeon Pea (*Cajanus cajan*)

Maryann Nkemakonam Anene¹ and Ernest Eguono Emojorho^{2*}

¹Department of Food Science and Technology, Nnamdi Azikiwe University, Awka, Nigeria

²Department of Food Science and Technology, University of Nigeria, Enugu State, Nigeria

Correspondence should be addressed to Ernest Eguono Emojorho, Department of Food Science and Technology, University of Nigeria, Enugu State, Nigeria

Received: October 07, 2023; Accepted: October 25, 2023; Published: November 01, 2023

ABSTRACT

This study looked at the idli's chemical, functional, physical, and sensory properties when prepared with rice, pigeon peas, and African yam beans. Dhals were made by washing, soaking whole pigeon pea and African yam bean (AYB) seeds in hot water (80°C for 30 minutes), hand-dehulling, and sun-drying them. For the processed dhals, black gram dhals (control), separate water was used to rinse and soak the rice for 5 hours at room temperature (29.2°C). The dhals were wet milled into a smooth batter separately from the rice, which was crushed to a coarse batter. Rice was combined 2:1 with dhals. The batters were combined, one gram of table salt was added, and the mixture was allowed to sit at room temperature (29.2°C) for 14 hours to ferment. Idli are created by pouring fermenting batter into moulds shaped like cups and heating them for 15 minutes. The chemical, functional, physical, and also the sensory properties of the flour and standard techniques were used to investigate the idli sample. The flour's protein content spanned the range of 7.88% to 26.27%. the carbohydrate content from 56.00% to 77.39%, and the ash content varied from 0.50% to 2.50% and 1.00% to 3.50%, respectively. Functional properties of the flours include bulk densities between 0.66 g/ml and 0.76 g/ml, swelling rates between 149.30% and 219.20%, and foaming rates between 7.69% and 34.62%. The moisture level of the idli samples varied from 63.00% to 68.50%, the protein content was between 4.31% and 6.79%, the fat content was between 0.10% and 2.50%, the ash content was between 0.50% and 2.00%, and the carbohydrate content was between 23.94% and 29.34%. The batter volume was from 109 ml to 122 ml, the texture was between 9.03 mm and 10.26 mm, the cooked weight was between 40.31 g and 49.91 g, the cooking time was between 7.03 minutes and 14.00 minutes, and the rise's diameter was between 7.16 cm and 7.70 cm. The idli samples all received favourable sensory evaluations. According to the sensory data rice, African yam beans, and pigeon peas can be combined to produce tasty idli.

KEYWORDS

Idli; Rice; Africa yam bean; Pigeon pea; Chemical; Functional

Citation: Maryann Nkemakonam Anene, Chemical Functional Physical and Sensory Properties of Flour and Idli Produced from Blends of Rice (*Oryza Sativa*), African Yam Bean (*Sphenostylis Sternocarpa*) and Pigeon Pea (*Cajanus Cajan*). Food Proc Nutr Sci 4(1): 39-50.

INTRODUCTION

Cereal-legume based foods are one of the primary sources of accessible dietary energy and minerals worldwide, according to [1]. Rice is one cereal grain that significantly influences human nutrition. Rice has high levels of cysteine and methionine but minimal quantities of several important amino acids, such as lysine. According to [2], rice cannot supply all the nutrients needed for a balanced diet on its own. Pulses, which are dietary supplements to a rice-based diet and include beans, black gram, pigeon pea, soybean, African yam bean, groundnuts, and lentils, complete the amino acid profile. To put it another way, rice-based dishes like masa, dosa, dhokla, and idli, among others, are made from a combination of rice and other legumes [3]. Idli is one of the cereal-legume based cuisines of India and is consumed three to four times a week for breakfast and dinner. It is a fermented steamed food that is unique to India and Sri Lanka. Idli is typically made with rice (*Oryza sativa*), a cereal, and black gram (*Phaseolus mungo*), a legume [4]. Idli is a food that naturally ferments; hence it is frequently fermented without the use of inoculum. This is so that the required microorganisms, including yeast and lactic acid bacteria, can be found naturally in the materials [5]. *Leuconostoc mesenteroides*, *Lactobacillus delbrueckii*, *Streptococcus faecalis*, and *Pediococcus cerevisiae* are a few examples of lactic acid bacteria. African yam beans are a locally underutilized legume, according to [6], despite not being widely consumed in Nigeria. It is widely acknowledged to have an exceptional amount of protein and most of the essential amino acids, including lysine and leucine. Pigeon peas, on the other hand, are a great source of carbohydrates, proteins, and vitamins. However, pulses like African yam bean (*Sphenostylis stenocarpa*) and Pigeon pea (*Cajanus cajan*), which are locally available and underutilized grain legumes in Nigeria [7], can be substituted in place of black gram in the production of idli due to their high nutritional values such as protein, carbohydrates, vitamins, and minerals. In order to avoid

this, it's important to look for cheap, high-quality protein sources, like pigeon peas and African yams.

Sources of Raw Materials

Pigeon pea and African yam bean grains were procured from Ogige market, Nsukka, Enugu State, Rice (Faro 44, Faro 52A and Faro 52B) was purchased from Abakiliki, Nigeria, and black gram (dhals) was purchased from Roban stores Enugu State, Nigeria.

Preparation of Pigeon Pea and African Yam Bean (AYB) Dhals

A Ohauz Adventurer Corporation Model AR2130 scale was used to weigh African yam bean and pigeon pea grain after they had been cleaned, sorted, and cleaned up. The seeds of the pigeon pea and African yam beans were soaked in boiling, 80 °C water for 30 minutes. After that, the hulls were painstakingly removed to make dhals with a mortar and pestle. The dhals were airtight jarred after being sun-dried and stored [8]. The process flow for making dhals out of pigeon peas and African yam beans (AYB) is shown in Figure 1.

Rice/black gram (control), Rice/African yam bean, and Rice/pigeon pea dhals




Figure 1: Processing of pigeon pea and African yam bean seeds into dhals.

The rice, black gram, African yam bean, and pigeon pea dhals were all ground separately in a hygienic atmosphere using a home blender. During milling, water additions ranging from 1.5 times to 2.2 times the rice and dhals' initial dry weight were made. The dhals were pounded into a finely smooth batter while the rice was then ground into a coarse batter. Each grain and legume were mixed with the rice batter. After receiving 1 g of table salt, the batters were combined. Then let the batters ferment for 14 hours at

29.1°C in a bowl that was covered. The pH, titratable acidity, and increase in batter volume were evaluated before and after fermentation. Equal amounts of batter samples were taken, placed in idli moulds stacked one on top of the other, and steamed for 7 minutes to 15 minutes in an idli pot. The idli pans/moulds were then oiled. Figure 2 depicts the production flow chart for idlis [9].




Figure 2: Production of African yam bean, pigeon pea, and black gram idlis.

Description of Idli Steaming Pot

The idli pot was modified and constructed in accordance with the standard idli pot, which consists of a chamber resembling a pot with a perforated cylindrical base and four moulds with cup-shaped depressions placed one on top of the other. Typically, the idli batter is put into the depressions that resemble cups, and they steam for 15 minutes. Figure 3 depicts the steaming pot for idlis. Figure 4 - Figure 6 show images of the black gram, African yam bean, and pigeon pea, respectively. In Figure 7, matured black gram idlis were displayed.



Figure 3: Idli steaming pots with mould pans stacked above the other.




Figure 4: Processed black gram.



Figure 5: African yam bean.



Figure 6: Pigeon pea dhals.



Figure 7: Developed black gram idlis.

Analytical Methods

The bulk density and foaming capacity were calculated using the techniques described in [10]. The “water absorb capacity” was assessed using a modified version of [11] methodology. Utilizing the techniques outlined by [12], the volume of batter was calculated. The swelling power was determined using the [13] method. A venier caliper that had been calibrated was used to measure the diameter of the formed idlis. Ohauz Adventurer Corporation Model AR2130 digital weighing balance was used to measure the

cooked weights of the developed idlis. Using a stopwatch Joerex digital timer no-4511-2, the cooking duration of the idli samples was calculated. Using the [14] method, the following values were determined: Moisture content, ash, crude protein, fiber, and fat. The difference was used as the method of total carbohydrate analysis.

Sensory Evaluation

The created idli samples were coded and given to a 20 persons panel from the University of Nigeria, Nsukka's department of Food Science and Technology for sensory evaluation. As illustrated in figure 8 and figure 9, the formulated Idli samples were delivered both alone (without accompaniments) and with accompaniments).



Figure 8: Samples of African yam bean, pigeon pea and black gram idlis.



Figure 9: Developed Idli samples with accompaniment.

Statistical Analysis

It was done using the completely randomized design (CRD). The statistical package for social sciences (SPSS) version 20 software was used to analyze the data using a one-way analysis of variance (ANOVA). Mean separation was by Duncan's New Multiple Range test. Significance was accepted at $p < 0.05$.

Experimental Design

The three different rice varieties Faro 44, 52A, and 52B were mixed with the three different legume dhals (African yam bean, pigeon pea, and black gram). The proportion of cereal to legume was 2:1. The analysis was completed using the experimental design shown below in Table 1.

Cereal (Rice)	Legumes (dhals)	Blend Ratio (g)	Sample Code
Faro 44(R1)	African Yam Bean	200:100	AR1
	Pigeon Pea	200:100	PR1
	Black Gram	200:100	BR1
Faro 52A(R2)	African Yam Bean	200:100	AR2
	Pigeon Pea	200:100	PR2
	Black Gram	200:100	BR2
Faro 52B(R3)	African Yam Bean	200:100	AR3
	Pigeon Pea	200:100	PR3
	Black Gram	200:100	BR3

Table 1: Design of experiment for blends of different rice cultivars, pigeon pea, African yam bean and black gram.

RESULTS AND DISCUSSION

Proximate Composition of Rice, African yam bean, Pigeon pea and Black gram Flours

The approximate proximate content of flours prepared from rice, African yam beans, pigeon peas, and black gram is shown in Table 2. The flours' moisture levels ranged from 10.46% to 12.60%. Faro 44 had the highest at 12.60%, which was higher than 10.69% and 11.77% observed for Faro 52A and Faro 52B respectively. There were statistically significant differences between the flours in terms of moisture content ($p < 0.05$). With these moisture values, it is anticipated that the flours will store well and be safe from bacterial action and mould growth [15]. These moisture content variations may be the result of varietal differences. The obtained moisture content values are thus low enough for the items, if packaged, to maintain proper shelf stability. All of the sample flours' protein levels varied considerably ($p < 0.05$) and ranged from 7.88% to 26.27%. The protein level of Faro 52B rice (9.85%) was lower than Faro 52A (8.93 %) and 7.88 % of Faro 44. Varietal variances may be the cause of these variations in protein concentration. Faro 52A protein contrasted favourably with the protein value of 8.35% as reported by [16], while Faro 44 protein was greater in content than (4.60%) as published

by [16]. According to [17], African yam beans were observed to contain between 24% and 28% protein. The samples of flour had a crude fiber level that ranged from 1.49% to 3.00%. Pigeon peas had the highest crude fiber content (3.00%) while Faro 52A rice had the lowest (1.49%). Crude fiber content of Faro 44 (1.86%) was higher than 1.50% recorded for Faro 52B. These rice flours' crude fiber contents were within the range (1.50% to 2.00%) found in [2] results. These differences in crude fiber content could be caused by varietal differences. Additionally, it was discovered that the black gram flour's fat level (1.4%) matched [18] findings. While [19] findings can be used to compare the fat content of pigeon pea flour. According to [15], the majority of legumes, including pigeon pea, have less than 3% fat content. The flours' individual ash levels ranged from 1.00% to 3.50%. Black gram flour was

discovered to have the greatest value of 3.5%, while Faro 44 and 52A rice flours had the lowest value of ash (1.00%). Pigeon pea, African yam bean, and Faro 44 and 52A did not vary significantly ($p > 0.05$). The fat contents found in the rice flours were comparable to [2] findings of 1.00% to 2.00%. The value of 2.2% that [20] reported as the ash content of AYB might be used to compare the two. The fact that black gram has such a high ash level indicates that its mineral richness is higher than other grains'. As a result, the samples' low ash concentration may have resulted from processing steps such drying, dehulling, and grinding [15]. The flours' varying carbohydrate levels, which were between 56.00% and 77.39%. Differences that were significant ($p < 0.05$) were seen. Faro 52A had the highest carbohydrate content (77.39%) while Faro 44 and Faro 52B had lower value of 76.16% and 74.88 % respectively.

Sample Flours	Moisture (%)	Protein (%)	Crude fibre (%)	Fat (%)	Ash (%)	Carbohydrate (%)
Faro 44	12.60 ^f ± 0.02	7.88 ^a ± 0.03	1.86 ^c ± 0.04	0.50 ^a ± 0.06	1.00 ^a ± 0.01	76.16 ^e ± 0.06
Faro52A	10.69 ^b ± 0.04	8.93 ^b ± 0.02	1.49 ^a ± 0.02	0.50 ^a ± 0.10	1.00 ^a ± 0.10	77.39 ^f ± 0.03
Faro52B	11.77 ^d ± 0.01	9.85 ^c ± 0.02	1.50 ^a ± 0.02	0.50 ^a ± 0.04	1.50 ^b ± 0.02	74.88 ^d ± 0.03
Pigeon pea	11.23 ^c ± 0.05	23.14 ^e ± 0.05	3.00 ^a ± 0.01	1.01 ^b ± 0.01	2.00 ^c ± 0.05	69.63 ^c ± 0.01
African yam bean	10.46 ^a ± 0.03	26.27 ^f ± 0.01	2.77 ^d ± 0.02	2.50 ^c ± 0.04	2.00 ^c ± 0.01	56.00 ^a ± 0.05
Black gram	12.00 ^e ± 0.02	19.72 ^d ± 0.05	1.60 ^b ± 0.05	1.00 ^b ± 0.03	3.50 ^d ± 0.10	62.18 ^b ± 0.01

Table 2: Proximate composition of different rice cultivars, pigeon pea, African yam bean and black gram flours. Values are means ± SD (n = 3). Samples with the same superscript in a row are not significantly different ($p > 0.05$)

Functional Properties of Rice, Pigeon pea, African yam bean and Black gram Flours

The results of the functional qualities of the flours from different types of rice, pigeon pea, African yam bean, and black gram are shown in Table 3. The flour samples' bulk densities ranged from 0.66 g/ml to 0.76 g/ml. Rice flours had the highest bulk density (0.76 g/ml), followed by that of pigeon pea flour (0.74 g/ml), and that of African yam

bean flour (0.66 g/ml) was the lowest. The bulk densities for rice flours were similar to [21] results which ranged from 0.511 g/cm³ to 0.878 g/cm³ for defatted and undefatted debittered orange seed flours. When compared favourably to the result (0.67 g/ml) achieved by [22], the African yam bean flour performed well. The results for pigeon pea flour (0.74 g/ml) were better than the results for wheat flour (0.71 g/ml) published by [23].

Samples	Bulk Density (g/ml)	Water Absorption Capacity (%)	Swelling Power (%)	Foaming Capacity (%)
Faro 44	0.76 ^e ± 0.00	134.10 ^a ± 0.15	149.30 ^a ± 1.43	7.69 ^a ± 2.00
Faro 52A	0.76 ^e ± 0.00	157.20 ^b ± 0.12	182.00 ^c ± 0.23	9.80 ^b ± 0.03
Faro52B	0.76 ^e ± 0.00	157.90 ^b ± 1.00	164.00 ^b ± 1.50	7.84 ^a ± 0.20
Pigeon pea	0.74 ^c ± 0.01	180.90 ^c ± 0.05	219.20 ^c ± 2.00	30.77 ^d ± 0.50
African yam bean	0.66 ^a ± 0.01	222.20 ^d ± 2.00	197.40 ^d ± 1.00	23.08 ^c ± 1.00
Black gram	0.71 ^b ± 0.02	262.00 ^e ± 0.20	216.90 ^e ± 1.70	34.62 ^e ± 1.00

Table 3: Functional Properties of flours from different rice cultivars, pigeon pea, African yam bean and black gram. Values are means ± SD (n = 3). Samples with the same superscript in a row are not significantly different ($p > 0.05$)

These flours had water absorption capabilities ranging from 134.10% to 260.00%. Faro 44 had the lowest result (134.00%), and the black gram flour sample had the greatest value (262.00%). Differences that were significant (p

> 0.05) were seen. These variations in water absorption capacity may be the result of different particle sizes.

Pigeon peas had a water absorption capacity that was comparable to the 188% and 189% values reported for them

and cowpeas, respectively, by [24]. The range of the flours' foaming capabilities was 7.69% to 34.62%. Differences that were significant ($p < 0.05$) were seen. The greatest amount of foam capacity was recorded in black gram flour (34.62%), followed by pigeon pea flour (30.77%), and Faro 44 rice flour had the lowest observed foam capacity (7.69%). The increased protein content of the grain and legume flours may be the cause of these elevated foam capacity.

The Proximate Composition of Idlis from Blends of Rice, Pigeon pea, African Yam Bean and Black Gram

Table 4 details the proximate composition of freshly prepared idlis manufactured from mixtures of rice, pigeon pea, African yam bean, and black gram. The idlis' moisture content rose from 63.00% to 68.50%. The control idli black gram/Faro 52B (BR3) had the lowest moisture content (63.00%), followed by the African yam bean/Faro 52B idli (AR3) and pigeon pea/Faro 52B idli (PR3). These high moisture contents could be as a result of soaking, the

amount of water added to the batter during product formulation and steaming process. These findings are comparable to [25] value (66.38%) for the moisture content of idlis. These outcomes also agree with those for cooked masa obtained by [26]. As a result of the product's high moisture content, it will therefore encourage bacterial activity and mould growth, which will unquestionably lower its shelf stability [15]. Idlis typically contain a lot of moisture and don't keep well for a long time. According to [27] report, a food's moisture content can be used to gauge its stability and susceptibility to microbial infection as well as to gauge its water activity. These increases in moisture occurred as a result of the idli products' formulation with water. The idlis' protein content increased from 4.31% to 6.70%. The highest value was found in the control idli black gram/Faro 52A (BR1) at 6.79%. This contrasted favourably with the figures [28] reported for conventional idli (7.0%). Pigeon pea/Faro 44 idli (PR1) had the next-highest protein level (5.47%), while pigeon pea/Faro 52B had the least protein value (4.31%).

Samples	Moisture (%)	Protein (%)	Crude fibre (%)	Fat (%)	Ash (%)	Carbohydrate (%)
AR1	66.50 ^g ± 0.05	4.60 ^b ± 0.02	0.40 ^b ± 0.01	1.00 ^e ± 0.04	1.00 ^b ± 0.01	26.50 ^e ± 0.01
BR1	65.00 ^d ± 0.02	5.25 ^d ± 0.04	0.35 ^b ± 0.04	1.00 ^e ± 0.01	2.00 ^d ± 0.05	26.40 ^d ± 0.03
PR1	64.00 ^b ± 0.03	5.47 ^e ± 0.03	0.52 ^e ± 0.02	2.00 ^d ± 0.03	1.00 ^b ± 0.05	27.01 ^f ± 0.02
AR2	65.25 ^c ± 0.01	5.25 ^d ± 0.04	0.48 ^d ± 0.01	2.50 ^e ± 0.03	2.00 ^d ± 0.01	24.52 ^b ± 0.05
BR2	64.50 ^c ± 0.04	6.79 ^f ± 0.03	0.23 ^a ± 0.03	1.00 ^e ± 0.04	1.00 ^b ± 0.03	26.48 ^e ± 0.05
PR2	65.50 ^f ± 0.01	4.31 ^a ± 0.01	0.66 ^f ± 0.05	1.00 ^e ± 0.02	1.00 ^b ± 0.01	27.53 ^g ± 0.01
AR3	68.50 ^h ± 0.03	5.25 ^d ± 0.05	0.71 ^f ± 0.02	0.10 ^a ± 0.03	1.50 ^c ± 0.04	23.94 ^a ± 0.01
BR3	63.00 ^a ± 0.02	4.82 ^c ± 0.02	0.84 ^g ± 0.05	1.00 ^e ± 0.05	1.00 ^b ± 0.02	29.34 ^h ± 0.03
PR3	68.50 ^h ± 0.01	5.25 ^d ± 0.04	0.43 ^{cd} ± 0.02	0.50 ^b ± 0.04	0.50 ^a ± 0.02	24.82 ^d ± 0.01

Table 4: Proximate composition of idlis from blends of different rice cultivars, pigeon pea, African yam bean and black gram.

*Values are means ± SD (n = 3). Samples with the same superscript in a row are not significantly different ($p > 0.05$). AR1=African yam bean/Faro 44, BR1=black gram/Faro 44, PR1=pigeon pea/Faro 44, AR2= African yam bean/Faro 52A, BR2=black gram/Faro 52A, PR2=pigeon pea/Faro 52A, AR3=African yam bean/Faro 52B, BR3=black gram/Faro 52B, PR3=pigeon pea/Faro 52B.

Therefore, at $p > 0.05$, there was no difference between the newly developed idlis made from African yam bean/Faro 52A (AR2), African yam bean/Faro 52B (AR3), and pigeon pea/Faro 52B (PR3) and the control idli made from black gram/Faro 44 (BR1).

The high moisture level of the idlis may be the cause of the drop in protein amounts of these samples. Pulses' protein content decreased as a result of dehulling [29]. The protein content of the idli samples was significantly reduced ($p < 0.05$) by steaming, which is consistent with observations

by [30] indicated heat treatment may reduce the relative compositions of protein. The protein content of fermented pearl millet decreased, according to [30]. The range of the crude fiber contents in the idli samples was 0.23% to 0.84%, and this difference was significant ($p > 0.05$). The black gram/Faro 52A idli (BR2) had the lowest value (0.23%), followed by the control idli (BR3), which had the greatest value (0.84%). The diluting effect brought on by a substantial amount of moisture content and the dehulling operations may be the cause of the low fiber content. Regarding their crude fiber content, the African yam bean and pigeon pea idli may be compared to the control idli. The

crude fiber contents of the other idli samples varied considerably ($p < 0.05$) from one another. These numbers matched up favourably with the crude fiber value (0.50%) discovered by [31]. The lowest crude fiber concentration was found in the control idli black gram/Faro 52A (BR2) at 0.23%. According to [32], fermentation significantly reduced fibre in fermented flours. However, it might be brought on by an increase in microflora populations and the metabolism's usage of fiber. These findings support the findings of [33], who found that boiling jackbeans reduced the amount of crude fiber. The idlis' fat content varied from 0.10% to 2.50%. African yam bean/Faro 52A idli (AR2) had the highest fat content (2.50%), whereas AR3 idli had the lowest fat content (0.10%). The high moisture level of the idlis and the steaming action, which caused lipids to be depleted, may have diluted the fat levels, while the vegetable oil used to lubricate the idli pans before to steaming may have contributed to the increased presence of fat. The amount of fat in PR3, BR3, PR2, BR2, AR1, and BR1 (control) was found to be consistent with the findings of [31] for control idli. It was not surprising to find that all of the idli samples had low fat concentrations because studies have demonstrated that legumes retain energy as carbohydrates rather than fat. All of the idli samples had significantly varying ash concentrations, ranging from 0.50% to 2.00% ($p < 0.05$). The highest value, 2.00%, was found in black gram/Faro 44 (BR1) and African yam bean/Faro 52A (AR2), while PR3 had the lowest value, 0.50%.

These low ash levels may be caused by the leaching of ash-forming minerals into soak water as well as during the dehulling and steaming operations. Idli samples suffered a substantial reduction due to the high moisture content, with reductions ranging from 23.94% to 29.34%. The PR3 idli had the lowest (23.94%) while the control idli (BR3) had the highest (29.34%). Idlis has lower carbohydrate levels than the figure (31.88%) reported by [31].

pH, Titratable Acidity and Rise in Batter Volumes of Idlis from Blends of Different Rice Cultivars, Pigeon Pea, African Yam Bean and Black Gram before and after 14 Hours Fermentation

Table 5 displays the pH titratable acidity and increase in batter volumes of idli batters made from mixtures of several rice kinds, pigeon pea, African yam bean, and black gram. A product's level of acidity or alkalinity is determined by its pH. The study's findings indicate that the pH of the idli batters varied from 6.20 to 6.60 at the beginning of fermentation. The greatest value (6.60) was found in the African yam bean/Faro 44 (AR1) and 52B (AR3) idli batters. The control idli black gram/Faro 52A (BR2) had the lowest pH of all the samples at 6.2. The black gram/Faro 44 (BR1) and black gram/Faro 52B (BR3) control idlis had a pH of 6.30, which was in good agreement with [34] observations that the pH of idli batters varied from 6.2 to 6.5 at the beginning of fermentation. The pH of the control idli batter was 6.32 when [35] first tested it; after 14 hours of fermentation, it had dropped to 4.53.

After 14 hours of spontaneous fermentation, the idli batters' pH values also varied from 4.50 to 4.80. The black gram/Faro 44, 52A, and 52B control idli batters were found to have the highest pH (4.80, 4.80, and 4.70, respectively). The newly developed idli batters (AR1, PR1, AR2, AR3, and PR3) with a pH of 4.50 showed lower values. These pH values were found to be consistent with those reported by [25] for control idli and after he added xylooligosaccharide to the batters, which varied from 4.69 to 4.57. According to [25], a reduction in pH (a rise in the amount of acid) would certainly encourage the growth of yeasts and, as a result, increase the creation of carbon dioxide, which leavens the batter. These legumes have a higher content of soluble nutrients to boost the growth of LAB after being soaked in water. According to [36], the purpose of LAB is to bring the batter's pH down to the ideal range (4.4 to 4.7) for yeast activity.

There was no significant ($p < 0.05$) difference in the titratable acidities of the idli samples before fermentation and ranged from 0.11% to 0.14%. The control idli batters BR1, BR2, and BR3 were found to have rapid fermentation and maximum values. However, the black gram/Faro 52B (BR3) control idli batter had the highest and lowest values, respectively. It has been suggested that the development of flavour and product stability is caused by acid generation. In the idli samples, an increase in the batter's acid level was seen as the natural fermentation process progressed. So, after 14 hours of fermentation, titratable acidity increased as the pH decreased. The microbial development, particularly of lactic bacteria, which produces lactic acid and reduce the pH and leavens the batter, may be to blame for the shift in pH and acid content [35]. Significant (p

< 0.05) increase was observed in the batter volumes of the idli samples while there was a noticeable change in the batter volume during natural fermentation. The amount of batter remaining after the 14-hours fermentation process varied between 109 ml to 122 ml. The pigeon pea idli batters showed the lowest values, with (PR1) having the lowest value of 109 ml. The biggest volume increase was seen in the control idli (BR1, BR2 and BR3). According to [37], the batter volume increased from 100 ml to 178.4 ml. As a result of the yeast's metabolic activity during natural fermentation, the volume may have increased for this reason [37]. It might also be ascribed to microbial development and production of enzymes that catalyze the breakdown of lipids, antinutritional substances, proteins, and carbohydrates [38].

Samples	pH before Fermentation	pH after Fermentation	TTA before Fermentation	TTA after Fermentation	Rise in Volume
			(%)	(%)	(ml)
AR1	6.6	4.5	0.12 ^a ± 0.00	0.50 ^{bc} ± 0.03	120 ^{de} ± 1.00
BR1	6.3	4.8	0.10 ^a ± 0.04	0.62 ^{de} ± 0.01	122 ^e ± 1.00
PR1	6.5	4.5	0.11 ^a ± 0.02	0.40 ^a ± 0.03	109 ^a ± 1.00
AR2	6.5	4.5	0.13 ^a ± 0.03	0.56 ^{cd} ± 0.03	115 ^c ± 0.57
BR2	6.2	4.8	0.13 ^a ± 0.00	0.64 ^e ± 0.00	118 ^d ± 1.00
PR2	6.4	4.6	0.13 ^a ± 0.01	0.42 ^a ± 0.08	110 ^{ab} ± 2.00
AR3	6.6	4.5	0.14 ^a ± 0.00	0.50 ^{bc} ± 0.05	114 ^c ± 2.51
BR3	6.3	4.7	0.12 ^a ± 0.02	0.68 ^e ± 0.00	118 ^d ± 2.00
PR3	6.5	4.5	0.11 ^a ± 0.02	0.46 ^{ab} ± 0.05	112 ^b ± 1.00

Table 5: pH, titratable acidity and rise in batter volume of idli batters from blends of different rice cultivars, pigeon pea, African yam bean and black gram before and after 14 hours fermentation.

*Values are means ± SD (n = 3). Samples with the same superscript in a row are not significantly different ($p > 0.05$). AR1=African yam bean/Faro 44, BR1=black gram/Faro 44, PR1=pigeon pea/Faro 44, AR2=African yam bean/Faro 52A, BR2=black gram/Faro 52A, PR2=pigeon pea/Faro 52A, AR3=African yam bean/Faro 52B, BR3=black gram/Faro 52B, PR3=pigeon pea/Faro 52B.

Physical Parameters of Developed Idlis from Blends of Different Rice Cultivars, Pigeon Pea, African Yam Bean and Black Gram

Table 6 lists the physical characteristics of developed idlis manufactured from mixtures of several rice kinds, pigeon pea, African yam bean, and black gram. The idli samples' diameters differed significantly from one another. The idli samples had diameters ranging from 7.16 cm to 7.7 cm. The diameter of the control idli (BR1) was 7.70 cm, while that of AR2 was 7.16 cm. Idlis, as reported by [39], are circular in shape and range in diameter from 7 cm to 10 cm, based on the dimension of the mold. The upper surface bulges, making the product thick in the center and taper toward the edges. This is consistent with the fact that idli often have a diameter of 7 cm to 10 cm. Due to the dependence on the

size of the steaming idli moulds, all of the idlis had a similar diameter. The cooked weights of the idli samples differed significantly ($p < 0.05$) from each other and varied from 40.31 g to 49.91 g. The cooked weights of the African yam bean idlis were higher than those of the control idlis, which had the lowest weights. The cooked weights of the African yam bean and pigeon pea idlis were found to be higher than those of the control idlis. This could be explained by fermentation, which made the black gram idli batter frothier and spongier than other batters.

The weight (50 g) that [28] were able to attain for a regular idli can be contrasted to these substantial weights of African yam bean idlis. The newly developed idli samples took between 7.03 minutes and 14.00 minutes to cook. There were significant differences ($p < 0.05$) between all the

samples. The pigeon pea idlis required more cooking time (13.02 minutes, 13.50 minutes, and 14.00 minutes), but the control idlis required less time (7.03 minutes, 7.04 minutes, and 7.07 minutes). The idli samples' textures had a range of 9.0 mm to 10.26 mm, and there were noticeable variances. The PR3 idli had a value of 9.03 mm, while the control idli (BR3) had a value of 10.26 mm. As a result, it is believed that the idlis with the greatest values have soft textures,

Sample s	Cooked Weights (g)	Cooking Time (min)	Diameter (cm)	Texture Measurements (mm)
AR1	49.91 ^d ± 0.58	12.12 ^b ± 0.02	7.43 ^{ab} ± 0.25	9.63 ^c ± 0.32
BR1	41.15 ^{ab} ± 0.73	7.04 ^a ± 0.03	7.70 ^b ± 0.20	10.10 ^d ± 0.10
PR1	44.42 ^{bc} ± 1.08	14.00 ^e ± 0.01	7.46 ^{ab} ± 0.28	9.16 ^{ab} ± 0.15
AR2	48.84 ^d ± 0.80	12.22 ^b ± 0.24	7.16 ^a ± 0.05	9.90 ^{cd} ± 0.36
BR2	40.57 ^a ± 0.56	7.03 ^a ± 0.02	7.43 ^{ab} ± 0.20	10.16 ^d ± 0.30
PR2	42.33 ^{abc} ± 1.89	13.02 ^c ± 0.03	7.36 ^{ab} ± 0.05	9.03 ^a ± 0.15
AR3	48.97 ^d ± 0.98	12.56 ^b ± 0.49	7.33 ^{ab} ± 0.25	10.20 ^d ± 0.20
BR3	40.31 ^a ± 0.94	7.07 ^a ± 0.03	7.66 ^b ± 0.15	10.26 ^d ± 0.25
PR3	44.12 ^{bc} ± 1.68	13.50 ^d ± 0.50	7.23 ^a ± 1.15	9.53 ^{bc} ± 0.25

Table 6: Physical parameters of developed idlis from blends of different rice cultivars, pigeon pea, African yam bean and black gram.

Values are means ± SD (n = 3). Samples with the same superscript in a row are not significantly different (p >0.05). AR1=African yam bean/Faro 44, BR1=black gram/Faro 44, PR1=pigeon pea/Faro 44, AR2=African yam bean/Faro 52A, BR2=black gram/Faro 52A, PR2=pigeon pea/Faro 52A, AR3=African yam bean/Faro 52B, BR3=black gram/Faro 52B, PR3=pigeon pea/Faro 52B.

Sensory Quality of Developed Idlis without Accompaniment from Blends of Different Rice Cultivars, Pigeon Pea, African Yam Bean and Black Gram

Table 7 and Table 8 illustrate the sensory characteristics of idlis with or without accompaniments. The findings demonstrated that when consumed with an accompaniment, the idli samples had appreciable ratings/scores for appearance, color, texture, flavour, taste, aftertaste, mouth feel, and overall acceptability. Comparing the pigeon pea/Faro 44 (PR1) and 52A (PR2) to the control black gram/Faro 44 (BR1) and 52A (BR2), there was no discernible difference in appearance (p >0.05). This indicates that consumers were also satisfied with the color of the freshly created idlis. The texture of the freshly created

while those with the lowest values are considered to have harsh textures. As a result, the control idlis were found to have a texture similar to that of the African yam bean/Faro 52B (AR3) and to be more easily pierced. According to [40], the idli produced after 14 hours of fermentation were found to be softer and more porous. The higher the values the softer the texture of the idlis while the lower the values the harder the texture of the idlis.

idlis was similar to that of the control idlis. As a result, there was no variation in texture between any of the idli samples that was significant (p >0.05). The control idli black gram/Faro 52A (BR2) was found to have the highest flavour qualities with a score of 7.30. This flavour reduction could be the result of volatile Flavors that were obtained through fermentation but were lost while steaming. Between the newly created samples PR1, AR2 and AR3 and the control idlis BR1, BR3, there was no discernible change (p >0.05). The taste, aftertaste, mouthfeel, and general acceptability of all the idli samples did not differ significantly (p >0.05). As a result, the tastes of all the samples were well received by the public. Thus, the newly formulated idli products were generally accepted by the consumers and possesses similar physical attributes with their control.

Samples	Appearance	Colour	Texture	Flavour	Taste	After taste	Mouthfeel	Overall Acceptability
AR1	7.25 ^a ± 1.11	5.90 ^{ab} ± 1.97	6.60 ^a ± 1.87	5.85 ^a ± 1.92	6.40 ^a ± 2.28	6.15 ^a ± 2.03	5.50 ^a ± 2.03	6.05 ^a ± 2.23
BR1	8.40 ^d ± 0.75	6.90 ^{bc} ± 1.74	6.35 ^a ± 2.27	6.65 ^{ab} ± 1.66	6.20 ^a ± 1.90	6.45 ^a ± 1.53	6.85 ^b ± 1.34	6.65 ^a ± 2.00
PR1	7.95 ^{cd} ± 0.75	6.90 ^{bc} ± 1.70	6.40 ^a ± 1.98	6.70 ^{ab} ± 2.10	6.60 ^a ± 1.95	5.80 ^a ± 2.04	6.05 ^{ab} ± 1.82	6.55 ^a ± 1.46
AR2	7.35 ^{abc} ± 0.74	5.90 ^{ab} ± 2.17	6.45 ^a ± 1.76	6.35 ^{ab} ± 2.20	6.25 ^a ± 1.74	6.00 ^a ± 1.58	5.60 ^{ab} ± 1.75	6.40 ^a ± 1.81
BR2	8.20 ^d ± 0.83	6.75 ^{bc} ± 1.91	6.65 ^a ± 2.00	7.30 ^b ± 1.30	6.10 ^a ± 2.02	5.85 ^a ± 1.87	6.25 ^{ab} ± 2.26	6.20 ^a ± 2.39
PR2	7.45 ^{abc} ± 0.94	6.70 ^{bc} ± 1.89	6.45 ^a ± 1.57	5.85 ^a ± 2.00	5.65 ^a ± 1.56	5.45 ^a ± 1.98	6.25 ^{ab} ± 1.68	6.25 ^a ± 1.86
AR3	7.90 ^{bcd} ± 0.78	5.25 ^a ± 2.51	6.30 ^a ± 1.41	6.60 ^{ab} ± 2.11	6.15 ^a ± 1.84	6.35 ^a ± 1.63	5.50 ^a ± 2.16	6.50 ^a ± 1.82
BR3	8.30 ^d ± 0.86	7.80 ^c ± 1.19	7.20 ^a ± 1.73	6.90 ^{ab} ± 1.55	6.50 ^a ± 1.96	6.25 ^a ± 1.55	6.70 ^{ab} ± 1.38	6.55 ^a ± 1.57
PR3	7.30 ^{ab} ± 1.12	6.50 ^{abc} ± 1.79	6.75 ^a ± 1.61	5.85 ^a ± 2.05	6.00 ^a ± 1.86	5.55 ^a ± 2.13	6.20 ^{ab} ± 1.57	6.50 ^a ± 1.27

Table 7: Sensory quality of developed idlis without accompaniment.

Values are means ± SD (n = 3). Samples with the same superscript in a row are not significantly different (p >0.05). AR1=African yam bean/Faro 44, BR1=black gram/Faro 44, PR1=pigeon pea/Faro 44, AR2= African yam bean/Faro 52A, BR2=black gram/Faro 52A, PR2=pigeon pea/Faro 52A, AR3=African yam bean/Faro 52B, BR3=black gram/Faro 52B, PR3=pigeon pea/Faro 52B.

Samples	Appearance	Colour	Texture	Flavour	Taste	Aftertaste	Mouthfeel	Overall Acceptability
AR1	7.45 ^a ± 1.35	7.10 ^{ab} ± 1.68	6.55 ^a ± 1.57	7.10 ^a ± 1.16	7.15 ^a ± 0.98	6.80 ^{ab} ± 1.36	6.55 ^a ± 1.57	7.00 ^{ab} ± 1.71
BR1	7.50 ^a ± 1.05	7.35 ^b ± 1.22	6.85 ^a ± 1.81	7.05 ^a ± 1.46	7.35 ^a ± 1.03	7.40 ^b ± 1.04	6.85 ^a ± 1.81	7.10 ^{ab} ± 0.91
PR1	7.25 ^a ± 1.48	6.95 ^{ab} ± 1.57	6.90 ^a ± 1.29	6.40 ^a ± 2.01	6.35 ^a ± 2.05	6.15 ^a ± 1.95	6.55 ^a ± 1.70	6.65 ^{ab} ± 1.34
AR2	7.65 ^a ± 0.98	6.55 ^{ab} ± 1.90	6.90 ^a ± 1.20	6.90 ^a ± 1.41	6.85 ^a ± 1.30	6.50 ^{ab} ± 1.57	6.85 ^a ± 1.13	7.10 ^{ab} ± 0.96
BR2	7.80 ^a ± 1.05	7.50 ^b ± 1.05	6.70 ^a ± 1.97	6.75 ^a ± 1.65	7.00 ^a ± 1.48	7.00 ^{ab} ± 1.12	7.30 ^a ± 1.41	7.10 ^a ± 1.16
PR2	7.30 ^a ± 0.97	7.15 ^{ab} ± 1.26	6.85 ^a ± 1.18	6.90 ^a ± 1.25	6.75 ^a ± 1.29	6.85 ^{ab} ± 1.26	7.10 ^a ± 1.07	6.25 ^a ± 1.77
AR3	7.65 ^a ± 1.13	6.15 ^a ± 1.69	6.65 ^a ± 1.34	6.65 ^a ± 1.42	6.80 ^a ± 1.32	6.85 ^{ab} ± 1.66	6.55 ^a ± 1.43	6.75 ^{ab} ± 1.25
BR3	7.75 ^a ± 1.11	7.45 ^b ± 1.53	7.05 ^a ± 1.23	6.90 ^a ± 2.10	6.85 ^a ± 1.75	6.65 ^{ab} ± 1.56	6.70 ^a ± 1.71	6.75 ^{ab} ± 1.51
PR3	7.25 ^a ± 0.91	7.30 ^b ± 1.26	6.85 ^a ± 1.08	6.75 ^a ± 1.80	7.05 ^a ± 1.14	7.50 ^b ± 1.14	7.00 ^a ± 1.41	7.35 ^b ± 1.22

Table 8: Sensory quality of developed idlis with accompaniment.

Values are means ± SD (n = 3). Samples with the same superscript in a row are not significantly different (p > 0.05). AR1=African yam bean/Faro 44, BR1=black gram/Faro 44, PR1=pigeon pea/Faro 44, AR2= African yam bean/Faro 52A, BR2=black gram/Faro 52A, PR2=pigeon pea/Faro 52A, AR3=African yam bean/Faro 52B, BR3=black gram/Faro 52B, PR3=pigeon pea/Faro 52B.

Sensory Quality of Developed Idlis with Accompaniment from Blends of Different Rice Cultivars, Pigeon Pea, African Yam Bean and Black Gram

Table 8 displays the findings of the sensory assessment and consumer acceptance of idlis manufactured from mixtures of several rice kinds, pigeon pea, African yam bean, and black gram. The results showed that the idli samples all had appreciable ratings/scores for appearance, colour, texture, flavour, taste, aftertaste, mouth feel and overall acceptability when consumed with an accompaniment. Overall acceptability scores ranged from 6.25 to 7.35. This demonstrates that the African yam bean/Faro 52A (AR2), black gram/Faro 44 (BR1), 52A (BR2), and pigeon pea/Faro 52B (PR3) idli samples received the highest mean scores of 7.10, 7.10, and 7.00, respectively, followed by the control idli samples. There were no significant variations between samples AR1, BR1, PR1, AR2, BR2, AR3, and BR3 (p > 0.05). This proves that consumers liked idlis more when they were eaten with a side dish.

CONCLUSION

This study has shown that mixes of rice, pigeon pea, and African yam bean can be used to make breakfast foods (idli) that are acceptable alternatives to black gram dhals. African yam beans, pigeon peas, and the Faro 44 and 52B rice types all produced idlis with protein level that was comparable to that of the black gram idlis (control). The results also showed that idlis made from blends of African yam beans with Faro 52A and 52B rice cultivars had higher values for ash content. African yam bean idli batters had greater volume to pigeon pea idli batters, in contrast to the control idli batters, according to the physical characteristics of the idlis. Consumers rated these idli goods highly and found them to be acceptable from a sensory and acceptance standpoint. Functional property outcomes have an impact on how flour is used and how many food products are developed. The flour's low bulk density makes it possible to formulate foods with less concern for retrogradation and has an impact on package design.

COMPETING INTERESTS

No conflict of interest or conflicting interests.

REFERENCES

1. Agrawal R, Rati ER, Vijayendra SVN, et al. (2000) Flavour profile of idli batter prepared from defined microbial starter cultures. *World Journal of Microbiology and Biotechnology* 16: 687-690.
2. Oko AO, Ugwu SI (2011) The proximate and mineral compositions of five major rice varieties in Abakaliki, South-Eastern Nigeria. *International Journal of Plant Physiology and Biochemistry* 3(2): 25-27.
3. Nkama I, Afodia KL, Amina AKJ (2011) Rice processing in Nigeria. Intellectual property and technology transfer office of the University of Maiduguri: 18-124.

4. Ghosh D, Chattopadhyay P (2011) Preparation of idli batter, its properties and nutritional improvement during fermentation. *Journal of Food Science and Technology* 48: 610-615.
5. Nagaraju VD, Manohar B (2000) Rheology and particle size changes during Idli fermentation. *Journal of Food Engineering* 43(3): 167-171.
6. Ene-Obong HN (1992) Nutritional evaluation, consumption pattern and processing of the African yam bean (*sphenostylis stenocarpa*) (Doctoral dissertation, PhD thesis, e University of Nigeria, Nsukka, Nigeria).
7. Azeke MA, Fretzdorff B, Buening-Pfaue H, et al. (2005) Nutritional value of African yambean (*Sphenostylis stenocarpa L*): Improvement by lactic acid fermentation. *Journal of the Science of Food and Agriculture* 85(6): 963-970.
8. Fasoyiro SB, Akande SR, Arowora KA, et al. (2010) Physico-chemical and sensory properties of pigeon pea (*Cajanus cajan*) flours. *African Journal of Food Science* 4(3): 120-126.
9. Steinkraus KH (1996) Handbook of indigenous fermented foods.(Revised and expanded) Marcel Dekker. Inc. New York.
10. Onwuka GI (2005) Food analysis and instrumentation: Theory and practice. Naphali prints.
11. Gandhi AP, Srivastava JAJFJ (2007) Studies on the production of protein isolates from defatted sesame seed (*Sesamum indicum*) flour and their nutritional profile. *ASEAN Food Journal* 14(3): 175.
12. Sridevi J, Halami PM, Vijayendra SVN (2010) Selection of starter cultures for idli batter fermentation and their effect on quality of idlis. *Journal of Food Science and Technology* 47: 557-563.
13. Leach HW, McCowen LD, Schoch TJ (1959) Structure of the starch granules. In: Swelling and solubility patterns of various starches. *Cereal Chemistry* 36: 534-544.
14. AOAC (2010) Official methods of analysis. Association of Official Analytical Chemists 18th (Edn.), Gaithersburg, Maryland, U.S.A.
15. Ihekoronye AI, Ngoddy PO (1985) Integrated food science and technology for the tropics. Macmillan: 10-77.
16. Badau MH, Ngozi C, Danbaba N (2013) Quality of garabia (A Nigerian traditional snack) from four varieties of rice as affected by the addition of cowpea. *Advance Journal of Food Science and Technology* 5(3): 249-254.
17. Adewale BD, Dumet DJ, Vroh-Bi I, et al. (2012) Morphological diversity analysis of African yam bean and prospects for utilization in germplasm conservation and breeding. *Genetic Resources and Crop Evolution* 59(5): 927-936.
18. Singh S, Singh HD, Sikka KC (1968) Distribution of nutrients in the anatomical parts of common Indian pulses. *Cereal Chemistry* 45: 13-18.
19. Faris DG, Saxena KB, Mazumdar S, et al. (1987) Vegetable pigeon pea: A promising crop for India: International Crop Research Institute for the Semi Arid Tropics, Patancheru, AP, India.
20. Oshodi AA, Ipinmoroti KO, Adeyeye EI (1997) Functional properties of some varieties of African yam bean (*Sphenostylis stenocarpa*) flour-III. *International Journal of Food Sciences and Nutrition* 48(4): 243-250.
21. Emojorho EE, Okonkwo TM (2022) Effect of debittering methods on the chemical and physiochemical properties of defatted and undefatted sweet orange seed flours. *Fruits* 77: 1-10.
22. Ojukwu M, Olawuni IA, Ibeabuchi C, et al. (2012) Physical, proximate and functional properties of 'Nsama'A local variety of African yam bean (*sphenostylis stenocarpa*) grown in southern states in Nigeria. *African Journal of Food Science and Technology* 3(10): 260-267.
23. Akubor PI, Badifu GI (2004) Chemical composition, functional properties and baking potential of African breadfruit kernel and wheat flour blends. *International Journal of Food Science & Technology* 39(2): 223-229.

24. Olalekan AJ, Bosede BF (2010) Comparative study on chemical composition and functional properties of three Nigerian legumes (jack beans, pigeon pea and cowpea). *Journal of Emerging Trends in Engineering and Applied Sciences* 1(1): 89-95.
25. Aachary AA, Gobinath D, Prapulla SG (2011) Short chain xylooligosaccharides: A potential prebiotic used to improve batter fermentation and its effect on the quality attributes of idli, a cereal-legume-based Indian traditional food. *International Journal of Food Science & Technology* 46(7): 1346-1355.
26. Nkama I (1993) Studies on improving the nutritional quality of masa. Traditional Nigerian fermented-cereal based food. Mysore: Central Food Technological Research Institute (CFTRI), United Nations University.
27. Frazier WC, Westhoff DC (1978) TMH edition. *Food microbiology*. New Delhi: Tata McGraw-Hill Publishing Company Ltd.
28. Nazni P, Shalini S (2010) Standardization and quality evaluation of idli prepared from pearl millet (*Pennisetum glaucum*). *International Journal of Current Research* 5: 84-87.
29. Singh U, Santosa BAS, Rao PV (1992) Effect of dehulling methods and physical characteristics of grains on dhal yield of pigeonpea (*Cajanus cajan L.*) genotypes. *International Journal Of Food Science & Technology* 29(6): 350-353.
30. El Hag ME, El Tinay AH, Yousif NE (2002) Effect of fermentation and dehulling on starch, total polyphenols, phytic acid content and in vitro protein digestibility of pearl millet. *Food Chemistry* 77(2): 193-196.
31. Dhanashree H, Danga SK, Patel MR, et al. (2015) A study on formulation, evaluation and analysis of soya-idli. *International Journal of Current Research and Academic Review* 3 (4): 145-155.
32. Nnam NM, Obiakor PN (2003) Effect of fermentation on the nutrient and antinutrient composition of baobab (*Adansonia digitata*) seeds and rice (*Oryza sativa*) grains. *Ecology of Food and Nutrition* 42(4-5): 265-277.
33. Udedibie ABI, Esonu BO, Unachukwu C, et al. (1996) Two-stage cooking as a method of improving the nutritive value of jackbean (*Canavalia ensiformis*) for broilers. *Nigerian Journal of Animal Production* 23(2): 107-110.
34. Sheela S Kowsalya S (2013) Microbial, physico-chemical and nutrient changes associated with Idli batter fermentation. *Global Research Analysis* 2: 74-75.
35. Rekha CR, Vijayalakshmi G (2011) Accelerated fermentation of 'idli' batter using soy residue okara. *Journal of Food Science and Technology* 48: 329-334.
36. Soni SK, Sandhu DK (1991) Role of yeast domination in Indian idli batter fermentation. *World Journal of Microbiology and Biotechnology* 7(4): 505-507.
37. Padmashree A, Semwal AD, Govindaraj T, et al. (2009) Effect of different drying methods on the stability of ready-to-reconstitute idlis. *International Journal of Advanced Research* 21: 2008-2010.
38. Rolle RS (1998) Enzyme applications for agro-processing in developing countries: An inventory of current and potential applications. *World Journal of Microbiology and Biotechnology* 14: 611-619.
39. Durgadevi M (2012) Enhancement of functional properties of Indian fermented food idli through incorporation of Moringa oleifera leave. *Journal of Scientific Research* 2(7): 213-215.
40. Balasubramanian S, Viswanathan R (2007) Properties of idli batter during its fermentation time. *Journal of Food Processing and Preservation* 31(1): 32-40.