

## Lipotransfer Strategies and Techniques to Achieve Successful Breast Reconstruction in the Radiated Breast

Kristina M Crawford<sup>1</sup> and Matthew R Endara<sup>2</sup>

<sup>1</sup>Resident Physician, General Surgery, Vanderbilt University Medical Center, Nashville, TN

<sup>2</sup>Maury Regional Medical Group, Columbia, TN; Clinical Assistant Professor of Surgery, Department of Surgery, Midwestern University, Downers Grove, IL

Correspondence should be addressed to Matthew R. Endara, [matt.endara@gmail.com](mailto:matt.endara@gmail.com)

Received: January 12, 2020; Accepted: January 26, 2020; Published: February 02, 2020

### **ABSTRACT**

Radiation therapy is frequently a critical component of breast cancer care but carries with it side effects particularly damaging to reconstructive efforts. The Autologous lipotransfer has the ability to improve radiated skin throughout the body due to the pluripotent stem cells and multiple growth factors transferred there in. The oncologic safety of lipotransfer to the breasts is demonstrated in the literature and is frequently considered an adjunctive procedure for improving aesthetics outcomes of breast reconstruction. Using lipotransfer as an integral rather than adjunctive step in the reconstructive process for breast cancer patients requiring radiation results in improved complication rates equivalent to non-radiated breasts; expanding options in these otherwise complicated cases.

### **Keywords**

Radiation therapy; Lipotransfer; Breast cancer; Reconstructive process

### **1. INTRODUCTION**

Reconstruction of the breasts following the surgical management of cancer is associated with improved quality of life, feelings of well-being and psychosocial development [1,2]. The objective of the reconstructive surgeon should be to offer options that facilitate these goals while minimizing potential complications. Use of radiation in the patient's oncologic care is a well-known and studied risk factor for increased complications and reconstructive failure [3]. Since the appropriate management of cancer frequently requires radiation treatment (RT) to improve survival and recurrence rates,

it is incumbent on the reconstructive surgeon to identify and implement strategies to compensate for this therapy. The traditional 'gold standard' treatment for this has been the transfer of well vascularized tissue in the form of a pedicled or free flap to reconstruct the resultant volume loss in the radiated breast [4]. Though effective, these procedures may not always be available or the patient's preferred option. Lack of access to properly trained reconstructive micro surgeons, inexperienced hospitals and a paucity of donor sites can seriously impact patients' ability to undergo these procedures. Patients

**Citation:** Kristina M Crawford, Lipotransfer Strategies and Techniques to Achieve Successful Breast Reconstruction in the Radiated Breast. Clin Surg J 3(3): 78-88.

© 2020 The Authors. Published by TRIDHA Scholars.

may also wish to avoid procedures with potential donor site complications, increased operative time, requisite inpatient admission or longer postoperative recovery times that can be associated with these more complex surgeries. Autologous lipotransfer is a relatively simple procedure that is being increasingly recognized as a strategy in the radiated patient with mounting evidence to support its use [5-13]. Though the ideal application of this technique remains debated in the literature, it is clearly becoming a critical and not simply adjunctive part of the reconstructive process in irradiated patients

## **2. USE OF RADIATION IN THE BREAST CANCER PATIENT**

The use of radiation treatments (RT) in breast cancer patients irrefutably improves survival and recurrence rates in lumpectomy and properly selected mastectomy patients [14-15]. Breast conservation as a treatment modality uses radiation as a necessary step. As such, the vast majority of these patients are subject to RT as part of their treatment. Indications for use in mastectomy patients are expanding as well with some centers offering it to as much as 70% of patients [16-17]. Indeed, meta-analysis by the Early Breast Cancer Trialists Collaborative Group (EBCTCG) have found improved rates of 10 year recurrence and 20 year mortality in doing so [14-15]. The number of patients who require reconstruction and have been or will be exposed to radiation is therefore increasing.

Irradiated tissues are associated with increased rates of surgical complications throughout the body [18-20]. This is especially true with radiated breast reconstructions as evidenced by higher rates of infection, capsular contracture, implant exposure and overall reconstructive failure [21-31]. Though still the most common form of reconstruction being performed today, staged expander to implant based reconstruction is especially sensitive to the unintended side effects of radiation. Complication rates in this patient population range from 38.9% to 68% [32-

33]. Consequently, some surgeons refuse to even offer this approach as they deem it a relative contraindication.

Patients desiring procedures to correct asymmetries, ptosis or macromastia following BCT are also at increased risk of complications. Several studies have looked directly at this patient population with generally mixed results [34]. Despite complications (e.g. delayed healing, prolonged edema, breast loss necessitating flap reconstruction), these procedures are believed to be safe but with careful patient selection. One study found a pooled complication rate of 50% with mastopexy or reduction in patients who had undergone BCT with RT [35]. Ultimately, an understanding of the harmful effect that radiation introduces to surrounding tissues is paramount to success when designing treatment strategies for reconstruction in these patients.

The mechanisms by which RT are so effective at shrinking tumor size and local recurrence are the same that cause the collateral side effects to local tissues. Radiation-induced tissue damage and the ensuing cellular and molecular response have been well-described in the literature [36-41]. The process occurs in three general phases: acute, latent, and late. The acute phase is thought to last from 0 to 6 months after exposure to radiation. This phase is characterized by damage to highly replicative cell lines through the initiation of cytokine cascades, creation of reactive oxygen species, and release of free radicals within the exposed cells. This property of RT is useful for causing apoptosis in cancer cells but is equally harmful to other proliferative cell lines such as basal keratinocytes and hair follicle stem cells. Damage to these regenerative cell lines results in the impairment of self-renewing abilities within the skin. Fibroblasts, endothelial cells, and epidermal cells within the radiation field are also affected resulting in the release of a variety of molecular signals. This leads to activation of the coagulation cascade as well as increased inflammation, tissue remodeling and epithelial regeneration. Finally,

blood vessels, especially smaller arterioles and capillaries, are affected during this phase. These vessels demonstrate increased permeability and thus tissue edema as well as the formation of fibrin plugs with resultant creation of local areas of ischemia.

The tissues proceed from the acute phase to a short latent period that is, yet, undefined but believed to begin approximately 6 months after treatment [42-43]. The late phase reactions occur next and can progress up to and beyond 20 years after initial exposure. Continued release of cytokines and growth factors results in prolonged fibroblast proliferation and progressive extracellular matrix deposition. Tissues becomes fibrotic with a decrease in vascular density. These factors lead to sites inhospitable to surgical interventions as they are stiff and have inadequate perfusion for healing. As such, these patients are frequently considered poor candidates for additional reconstructive procedures. Several studies have identified expander to implant surgeries to be particularly susceptible to these negative effects [3,4,21]. Patients desiring some form of breast reshaping after breast conservation therapy are equally approached with caution.

Strategies that taken into account, these harmful effects of radiation have been met with some success [15,17,44-51]. This includes delaying the expander to implant exchange procedure for 6 months to allow for completion of the acute phase, using a counter incision in the IMF thereby avoiding the more heavily radiated mastectomy incision line, and use of autologous lipotransfer to physiologically reverse the harmful effects.

### **3. AUTOLOGOUS LIPOTRANSFER TO REGENERATE RADIATED TISSUES**

Though used for over 100 years to increase tissue bulk for cosmetic effect throughout the body, autologous lipotransfer is now being seen as a particularly useful technique for treating radiodermatitis [52-53]. The reason

it is so effective in this regard appears to be from the multipotent adipose derived stem cells, adipose derived regenerative cells transferred and miscellaneous components of the stromal vascular fraction of the graft. The exact mechanism by which these cell lines are capable of reversing the harmful effects of radiation is as of yet unknown. It does appear that within radiated tissue the ASC is important as it is capable of thriving and even proliferating in that ischemic environment [54]. Suspected mechanisms for the ASCs to promote reversal of radiodermatitis are their ability to differentiate into lost cell types and to release paracrine signals with proangiogenic and anti-fibrotic effects.

Another factor that may contribute to the proangiogenic effect seen with lipotransfer into irradiated tissues is the inclusion of additional vessel forming elements with the SVF [55]. These include endothelial cells, pericytes, smooth muscle cells and their progenitors capable of forming vascular cells and blood vessels. Experimental models transferring human fat to irradiated murine tissue has supported these findings. Indeed, the grafted tissue was found to have decreased dermal thickness, reduction in collagen content, increase in vascular density and overall improved fat graft retention.

Clinically the beneficial effect of fat grafting in radiated patients has been demonstrated in several studies [8-10]. This technique for ameliorating radiodermatitis of the breast is changing the way that reconstructive surgeons are approaching breasts cancer patients. Initial concern over the potential for cancer activation limited the use of autologous lipotransfer in the breast. Multiple clinical studies, meta-analysis and systematic reviews have failed to provide evidence to support this concern. Consequently, lipofilling had been used with increasing popularity in breast reconstruction but typically as an adjunctive step to improve the final cosmetic result [56]. Increased recognition of the positive effects on radiation tissue, however, is resulting in the development of

treatment protocols that incorporate it as an integral part of reconstructing these radiated patients [5,7].


#### **4. USE OF AUTOLOGOUS LIPOTRANSFER IN RECONSTRUCTION OF THE RADIATED BREAST**

Initial experience with lipotransfer in the radiated breast focused on using it to revive and prime post mastectomy skin flaps either after the completion of reconstruction or prior to attempting it [6,11-13]. These strategies were important for demonstrating efficacy in improving complication and failure rates but limited in the cosmetic results they were able to obtain and delaying the overall time course.

Building on this, Ribuffo et al. [5] presented 32 patients who underwent modified radical mastectomy followed by RT. Patients were reconstructed in an immediate fashion at the time of mastectomy with placement of tissue expanders in a submuscular plane. Half of the patients underwent between 1-2 separate autologous lipotransfer procedures as early as 6 weeks after completion of radiotherapy before expander to implant exchange. They reported a 0% complication rate in their treatment arm and a 43% in the control group. Introducing lipotransfer as a separate but necessary part of their protocol was unique as it became a formal part of their protocol for success.

Work by Serra-Renom et al. [7] confirmed the utility of lipotransfer in 65 of their mastectomized irradiated patients by incorporating serial fat grafting into their protocol. These patients underwent multiple fat grafting procedures including before and at the time of expander to implant exchange. They found excellent clinical results with their technique. This study was limited as patients were not demonstrating significant acute effects of radiation in the form of radiodermatitis and thus the severity of damage to the tissues was in question.

Our 3-stage lipo-approach to mastectomized irradiated patients is modeled on these previous studies and additional best available evidence for mitigating radiotoxicity. The hallmarks of our algorithm include: use of ADM, maintenance of the expander in a fully inflated position during radiation, delay of the expander-to-implant procedure for at least 6 months after radiotherapy completion, use of a counter-incision at the IMF in cases of skin-sparing mastectomy (SSM), and performance of a separate surgery whereby autologous fat was transferred to the radiated breast prior to the final exchange. Comparing radiated breasts to our general non-radiated population as well as within for patients who had bilateral mastectomy whereby one breast was radiated and one non-radiated revealed equivalent complication rates ( $p = 0.387$  and  $p = 1$  respectively) (Figure 1).



**Figure 1A:** Prior to bilateral mastectomy and 6 weeks after attempted lumpectomy with positive margins.



**Figure 1B:** Ten weeks status-post bilateral mastectomy with immediate tissue expander placement and halfway through radiotherapy regime, with severe radiodermatitis.



**Figure 1C:** Twenty-two months following expander-to-implant exchange.

A subset of patients who have undergone breast conservation therapy with lumpectomy followed by radiation desire either mastopexy or reduction to improve their postoperative appearance as well as improve symmetry between their treated and untreated breast. Current recommendations are to limit offering these surgeries to carefully selected patients as long after radiation as possible. Use of fat grafting to ‘prime’ the skin envelope as a separate procedure prior to attempting a reduction or lift is an alternative strategy that has been successfully utilized in our practice (non published) with reproducible and reliable results (Figure 2).



**Figure 2A:** One year following completion of BCT with severe asymmetry and bilateral ptosis



**Figure 2B:** Following completion of fat grating to radiated breast, small reduction to non-radiated breast.




**Figure 2C:** Six months following completion of mastopexy to radiated breast.

#### *4.1 Three Step Approach in the Mastectomy Patient requiring post mastectomy radiation*

In radiation naïve patients undergoing mastectomy preoperative consultation includes a discussion of the 3-stage approach to implant-based reconstruction as well as the use of autologous flaps should radiation be in question or required. Please see Figure 3. Inherent in this approach is the use of flaps as a salvage procedure should implant failure arise. Patients who have already undergone radiation, such as previous breast conservation therapy patients now suffering a recurrence, are not candidates for this approach and some form of autologous flap transfer is recommended.

Regardless of whether a patient would prefer flap or implant-based reconstruction, we prefer to proceed in a delayed immediate fashion. Placement of a tissue expander at the time of mastectomy is therefore essential in this technique precluding direct to implant or immediate flap reconstruction. Though evidence for use

of this technique applies to partial submuscular expander placement, it is currently being used for prepectoral reconstructions as well [57]. In either case the use of acellular dermal matrix is considered an important part of the expander placement. Placement of this matrix creates a plane for ease of lipotransfer and has proven itself to be radioprotective [58].



**Figure 3:** Three stage treatment protocol to for complete breast reconstruction following mastectomy.

The patient undergoes serial expansions starting two weeks after the index procedure. The breast to be radiated is expanded to patient preference or expander limit. Maintaining the expander in an inflated position is important for success and has been shown to be preferable from a radiation standpoint as well. In those who have undergone bilateral mastectomy the non-radiated side is deflated prior to radiation to protect the heart and lungs. The patient is monitored through their RT with a clinic visit at the halfway mark and following completion of treatment. One week after completion of radiation the non-radiated breast is then easily re-inflated.

The patient is then subject to a 3-month waiting period prior to their next procedure: autologous lipotransfer to the radiated breast. At this point the patient is taken back to the operating room for whole breast fat grafting. This procedure is performed by utilizing a super wet technique for liposuction into a revolved fat transfer harvest system. Prior to injection of the fat pre-tunneling is performed within the subcutaneous space. The correct space is identified under direct visualization through a 1cm incision within the mastectomy scar. Avoidance of using sharp tipped instruments is important during this step to avoid rupturing the expander. Scar tissue bands that may block the ability to uniformly inject the fat are severed using a riveted fat harvest cannula with a saw type motion. The expander is then deflated by 60 ml-100 ml to make room for transfer of the fat. It is important to avoid over deflation as the fat still requires a flat plane to be placed in a string of pearls fashion.

The fat is then processed with three washes of warm lactated ringers. It is transferred from the revolve to a 60 ml syringe then into 3 ml syringes for transfer. The majority of the fat is injected through the incision. If additional access sites are needed to optimize the angle of delivery, they can be made with a 16-gauge needle while tenting the skin from the inside with a fat grafting cannula to protect the device. Constant motion of the syringe while injecting the fat is critical to avoid clumping of the fat graft and subsequent poor take. Enough fat is injected to fill the space by injecting at least as much as the fluid removed but not to the point of skin discoloration or creation of an overly taut skin envelope.

Following completion of this step, the patient is monitored closely with weekly follow up visits for the 3 weeks after surgery. If additional expansion is desired this can be attempted at 1-month post fat grating. The patient is subject to another 3-month waiting period before their final surgery. This is timed to optimize the



chances the lipoaspirate will positively affect the acute phase of radiation injury while performing the surgery during the latent phase.



**Figure 4A:** Newly diagnosed right breast cancer prior to bilateral mastectomy.



**Figure 4B:** Five months after bilateral mastectomy with immediate tissue expander placement and three months after radiotherapy completion, photo taken on day of lipotransfer surgery with markings for fat harvest from abdomen.

The patient is taken back to the operating room once more for removal of the expander and placement of the final implant. In cases of skin sparing mastectomy, a counter incision is utilized within the inframammary fold. Patients who have undergone a nipple sparing mastectomy using an IMF incision are accessed by extending the incision laterally. Any requisite capsular modifications are able to be performed at this time to not only enhance the final shape of the reconstructed breast

but also address any contraction that had occurred as a result of radiation. The implant is introduced using a touch free technique. Closure is performed in layers using buried, monofilament dissolvable suture (Figure 4).



**Figure 4C:** Seven months following lipotransfer to right breast and four months post expander-to-implant exchange.

#### *4.2 Two staged approach to BCT patients desiring oncoplastic mastopexy or reduction after completion of RT*

Eligible patients typically present with severe asymmetry following the completion of their breast conservation therapy combined with ptosis and/or macromastia. These patients are counseled on the risks of operating in a previously radiated surgical field and the benefit of priming the tissues with autologous lipotransfer before doing so. A two-step approach is offered with the first procedure consisting of a lift/reduction of the non-radiated breast and fat grafting to the radiated breast. The mastopexy/reduction of the radiated side is performed at least 3 months later.

The fat grafting technique is similar as for mastectomy patients with the preferred plane of injection remaining in the subcutaneous space. Care is taken not to inject directly into the breast tissues for multiple reasons. The subcutaneous plane is the target for transfer as we are trying to reverse the negative radiation effects on those regenerative cell lines that if damaged will lead to

increased risk with the final breast shaping procedure. Injection into breast tissue would not accomplish this goal and though not proven by data would be more concerning from an oncologic perspective. Multiple patients have been operated on in this manner by the primary author with the largest complication being persistent asymmetry that was still improved from before surgery. The technique has allowed for reduction in the selectivity for offering breast reshaping in previously radiated fields as has been recommended by previous authors.

## **5. CONCLUSIONS**

Recognition of the regenerative properties of lipoaspirate in radiated fields is leading to simplification of breast

reconstruction in this otherwise complicated patient population. The optimal strategy for this requires knowledge of the harmful effects of radiation, their time course and the biomolecular pathways by which lipoaspirate reverses them. Though not yet fully understood it is clear that applying autologous lipotransfer can significantly improve breast reconstruction outcomes in irradiated patients. The ease of these fat grafting procedures along with their use and application throughout the body makes them very well known to most plastic surgeons and may be leading to a paradigm shift in approaching the radiated breast. This has the potential to improve options and access to reconstruction for this ever-growing patient group.

## **References**

1. Fanakidou I, Zyga S, Alikari V, et al. (2018) Mental health, loneliness, and illness perception outcomes in quality of life among young breast cancer patients after mastectomy: The role of breast reconstruction. *Quality of Life Research* 27(2): 539-543.
2. Sisco M, Johnson DB, Wang C, et al. (2015) The quality-of-life benefits of breast reconstruction do not diminish with age. *Journal of Surgical Oncology* 111(6): 663-668.
3. Mericli AF, Sharabi SE (2019) Breast implants and radiation. *Seminars in plastic surgery*, MD Anderson cancer Center, the University of Texas, USA.
4. See MS-F, Farhadi J (2018) Radiation therapy and immediate breast reconstruction: Novel approaches and evidence base for radiation effects on the reconstructed breast. *Clinics in Plastic Surgery* 45(1): 13-24.
5. Ribuffo D, Atzeni M, Guerra M, et al. (2013) Treatment of irradiated expanders: protective lipofilling allows immediate prosthetic breast reconstruction in the setting of postoperative radiotherapy. *Aesthetic Plastic Surgery* 37(6): 1146-1152.
6. Rigotti G, Marchi A, Galiè M, et al. (2007) Clinical treatment of radiotherapy tissue damage by lipoaspirate transplant: a healing process mediated by adipose-derived adult stem cells. *Plastic and Reconstructive Surgery* 119(5): 1409-1422 .
7. Serra-Renom JM, Muñoz-Olmo JL, Serra-Mestre JM (2010) Fat grafting in postmastectomy breast reconstruction with expanders and prostheses in patients who have received radiotherapy: formation of new subcutaneous tissue. *Plastic and Reconstructive Surgery* 125(1): 12-18.
8. Jackson IT, Simman R, Tholen R, et al. (2001) A successful long-term method of fat grafting: Recontouring of a large subcutaneous postradiation thigh defect with autologous fat transplantation. *Aesthetic Plastic Surgery* 25(3): 165-169.
9. Kim SS, Kawamoto HK, Kohan E, et al. (2010) Reconstruction of the irradiated orbit with autogenous fat grafting for improved ocular implant. *Plastic and Reconstructive Surgery* 126(1): 213-220.
10. Phulpin B, Gangloff P, Tran N, et al. (2009) Rehabilitation of irradiated head and neck tissues by autologous fat transplantation. *Plastic and Reconstructive Surgery* 123(4): 1187-1197.



11. Missana MC, Laurent I, Barreau L, et al. (2007) Autologous fat transfer in reconstructive breast surgery: indications, technique and results. *European Journal of Surgical Oncology* 33(6): 685-690.
12. Panettiere P, Marchetti L, Accorsi D (2009) The serial free fat transfer in irradiated prosthetic breast reconstructions. *Aesthetic Plastic Surgery* 33(5): 695-700.
13. Salgarello M, Visconti G, Barone-Adesi L (2012) Fat grafting and breast reconstruction with implant: another option for irradiated breast cancer patients. *Plastic and Reconstructive Surgery* 129(2): 317-329.
14. Clarke M, Collins R, Darby S, et al. (2005) Effects of radiotherapy and of differences in the extent of surgery for early breast cancer on local recurrence and 15-year survival: An overview of the randomized trials. *Lancet* 366(9503): 2087-2106.
15. McGale P, Taylor C, Correa C, et al. (2014) Effect of radiotherapy after mastectomy and axillary surgery on 10-year recurrence and 20-year breast cancer mortality: meta-analysis of individual patient data for 8135 women in 22 randomized trials. *Lancet* 383(9935): 2127-2135.
16. American Cancer Society (2018) *Breast Cancer Facts & Figures 2017-2018*. Atlanta: American Cancer Society.
17. Delaney G, Barton M, Jacob S (2003) Estimation of an optimal radiotherapy utilization rate for breast carcinoma: a review of the evidence. *Cancer* 98(9): 1977-1986.
18. Robinson DW (1955) The hazards of surgery in irradiated tissue. *The Archives of Surgery* 71(3): 410-418.
19. Robinson DW (1975) Surgical problems in the excision and repair of radiated tissue. *Plastic and Reconstructive Surgery* 55(1): 41-49
20. Rudolph R (1982) Complications of surgery for radiotherapy skin damage. *Plastic and Reconstructive Surgery* 70(2): 179-185.
21. Cordeiro PG, Pusic AL, Disa JJ, et al. (2004) Irradiation after immediate tissue expander/implant breast reconstruction: outcomes, complications, aesthetic results, and satisfaction among 156 patients. *Plastic and Reconstructive Surgery* 113(3): 877-881.
22. Jhaveri JD, Rush SC, Kostroff K, et al. (2008) Clinical outcomes of postmastectomy radiation therapy after immediate breast reconstruction. *International Journal of Radiation Oncology, Biology, Physics* 72(3): 859-865.
23. Tallet AV, Salem N, Moutardier V, et al. (2003) Radiotherapy and immediate two-stage breast reconstruction with a tissue expander and implant: complications and esthetic results. *International Journal of Radiation Oncology, Biology Physics* 57(1):136-142.
24. Krueger EA, Wilkins EG, Strawderman M, et al. (2001) Complications and patient satisfaction following expander/implant breast reconstruction with and without radiotherapy. *International Journal of Radiation Oncology, Biology, Physics* 49(3): 713-721.
25. Kronowitz SJ, Hunt KK, Kuerer HM, et al. (2004) Delayed-immediate breast reconstruction. *Plastic and Reconstructive Surgery* 113(6): 1617-1628.
26. Spear SL, Majidian A (1998) Immediate breast reconstruction in two stages using textured, integrated-valve tissue expanders and breast implants: a retrospective review of 171 consecutive breast reconstructions from 1989 to 1996. *Plastic and Reconstructive Surgery* 101(1): 53-63.
27. Alderman AK, Wilkins EG, Kim HM, et al. (2002) Complications in postmastectomy breast reconstruction: two-year results of the Michigan Breast Reconstruction Outcome Study. *Plastic and Reconstructive Surgery* 109(7): 2265-2274.

28. Ricci JA, Epstein S, Momoh AO, et al. (2017) A meta-analysis of implant-based breast reconstruction and timing of adjuvant radiation therapy. *The Journal of Surgical Research* 218: 108-116.
29. Ascherman JA, Hanasono MM, Newman MI, et al. (2006) Implant reconstruction in breast cancer patients treated with radiation therapy. *Plastic and Reconstructive Surgery* 117(2): 359-365.
30. Anker CJ, Hymas RV, Ahluwalia R, et al. (2015) The Effect of radiation on complication rates and patient satisfaction in breast reconstruction using temporary tissue expanders and permanent implants. *Breast Journal* 21(3): 233-240.
31. Sullivan SR, Fletcher DR, Isom CD, et al. (2008) True incidence of all complications following immediate and delayed breast reconstruction. *Plastic and Reconstructive Surgery* 122(1): 19-28.
32. Poppe MM, Agarwal JP (2017) Breast reconstruction with postmastectomy radiation: Choices and tradeoffs. *Journal of Clinical Oncology* 35(22): 2467-2470.
33. Jagsi R, Momoh AO, Qi J, et al. (2018) Impact of radiotherapy on complications and patient-reported outcomes after breast reconstruction. *Journal of The National Cancer Institute* 110(2).
34. Barnea Y, Bracha G, Arad E, et al. (2019) Breast reduction and mastopexy for repair of asymmetry after breast conservation therapy: Lessons learned. *Aesthetic plastic surgery* 43(3): 600-607.
35. Spear SL, Rao SS, Patel KM, et al. (2014) Reduction mammoplasty and mastopexy in previously irradiated breasts. *Aesthetic Surgery Journal* 34(1): 74-78.
36. Hymes SR, Strom EA, Fife C (2006) Radiation dermatitis: Clinical presentation, pathophysiology, and treatment 2006. *Journal of the American Academy of Dermatology* 54(1): 28-46.
37. Hauer-Jensen M, Fink LM, Wang J (2004) Radiation injury and the protein C pathway. *Critical Care Medicine* 32(5): S325-S330.
38. López E, Guerrero R, Núñez MI, et al. (2005) Early and late skin reactions to radiotherapy for breast cancer and their correlation with radiation-induced DNA damage in lymphocytes. *Breast Cancer Research* 7(5): R690-R698.
39. Yarnold J, Brotons M-CV (2010) Pathogenetic mechanisms in radiation fibrosis. *Radiotherapy & Oncology* 97(1): 149-161.
40. Martin M, Lefaix J-L, Delanian S (2000) TGF- $\beta$ 1 and radiation fibrosis: a master switch and a specific therapeutic target? *International Journal of Radiation Oncology, Biology, Physics* 47(2): 277-290.
41. Wei J, Wang B, Wang H, et al. (2019) Radiation-induced normal tissue damage: Oxidative stress and epigenetic mechanisms. *Oxidative Medicine and Cellular Longevity*.
42. Bentzen SM, Thames HD, Overgaard M (1989) Latent-time estimation for late cutaneous and subcutaneous radiation reactions in a single-follow-up clinical study. *Radiotherapy and Oncology* 15(3): 267-274.
43. Arcangeli G, Friedman M, Paoluzi R (1974) A quantitative study of late radiation effect on normal skin and subcutaneous tissues in human beings. *The British Journal of Radiology* 47(553): 44-50.
44. Nahabedian MY (2013) Minimizing incisional dehiscence following 2-stage prosthetic breast reconstruction in the setting of radiation therapy. *Gland Surgery* 2(3): 133-136.
45. Fowble B, Park C, Wang F, et al. (2015) Rates of reconstruction failure in patients undergoing immediate reconstruction with tissue expanders and/or implants and postmastectomy radiation therapy. *International Journal of Radiation Oncology, Biology, Physics* 92(3): 634-641.
46. Kronowitz SJ (2015) State of the art and science in postmastectomy breast reconstruction. *Plastic and Reconstructive Surgery* 135(4): 755e-771e.

47. Percec I, Bucky LP (2008) Successful prosthetic breast reconstruction after radiation therapy. *Annals of Plastic Surgery* 60(5): 527-531.
48. Kronowitz SJ, Robb GL (2009) Radiation therapy and breast reconstruction: A critical review of the literature. *Plastic and Reconstructive Surgery* 124(2): 395-408.
49. Kronowitz SJ (2012) Current status of implant-based breast reconstruction in patients receiving postmastectomy radiation therapy. *Plastic and Reconstructive Surgery* 130(4): 513e-523e.
50. Trovo M, Durofil E, Polesel J, et al. (2012) Locoregional failure in early-stage breast cancer patients treated with radical mastectomy and adjuvant systemic therapy: which patients benefit from postmastectomy irradiation? *International Journal of Radiation Oncology, Biology, Physics* 83(2): e153-e157.
51. Recht A, Edge SB, Solin LJ, et al. (2001) Postmastectomy radiotherapy: clinical practice guidelines of the American Society of Clinical Oncology. *Journal of Clinical Oncology* 19(5): 1539-1569.
52. Simonacci F, Bertozzi N, Grieco MP, et al. (2017) Procedure, applications, and outcomes of autologous fat grafting. *Annals of Medicine and Surgery* 20: 49-60.
53. Borrelli MR, Shen AH, Lee GK, et al. (2019) Radiation-Induced skin fibrosis: Pathogenesis, current treatment options, and emerging therapeutics. *Annals of Plastic Surgery* 83(4S): S59-S64.
54. Kumar R, Griffin M, Adigbli G, (2016) Lipotransfer for radiation-induced skin fibrosis. *British Journal of Surgery* 103(8): 950-961.
55. Hong KY, Yim S, Kim HJ, et al. (2018) The fate of the adipose-derived stromal cells during angiogenesis and adipogenesis after cell-assisted lipotransfer. *Plastic and Reconstructive Surgery* 141(2): 365-375.
56. Zheng DN, Li QF, Lei H, et al. (2008) Autologous fat grafting to the breast for cosmetic enhancement: Experience in 66 patients with long-term follow up. *Journal of Plastic and Reconstructive Aesthetic Surgery* 61(7): 792-798.
57. Sbitany H, Piper M, Lentz R (2017) Prepectoral breast reconstruction: A safe alternative to submuscular prosthetic reconstruction following nipple-sparing mastectomy. *Plastic and Reconstructive Surgery* 140(3): 432-443.
58. Borrelli MR, Patel RA, Sokol J, et al. (2019) Fat chance: The rejuvenation of irradiated skin. *Plastic and Reconstructive Surgery - Global Open* 7(2): e2092.