

CLINICAL RESEARCH

# Nurses Practice and Associated Factors towards Hemodialysis Care in Addis Ababa, Ethiopia

Temesgen Legesse<sup>1</sup>, Amanuel Fanta<sup>2\*</sup>, Teshale Belayneh<sup>3</sup> and Tamrat Legesse<sup>4</sup>

<sup>1</sup>Addis Ababa University, Black Lion Hospital, Ethiopia

<sup>2</sup>Department of Nursing, Hawassa College of Health Sciences, Ethiopia

<sup>3</sup>Department of Public Health, Hawassa College of Health Sciences, Ethiopia

<sup>4</sup>Department of Health sciences, Dilla University, Ethiopia

Correspondence should be addressed to Amanuel Fanta, Department of Nursing, Hawassa College of Health Sciences, Ethiopia

Received: 23 February 2023; Accepted: 05 March 2023; Published: 12 March 2023

Copyright © Temesgen Legesse. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## **ABSTRACT**

### **BACKGROUND**

Patients on hemodialysis care are at a greater risk of developing infectious diseases that leads to life treating complications. The deficiency in practice of nurses when caring for patients who are in hemodialysis has a negative effect on the improvement of the patient's condition. This study was conducted to determine nurse's practices and associated factors towards dialysis care.

### **METHODS**

An institution based cross-sectional study was employed in this study. A total of 62 nurses giving hemodialysis care were included. Statistical analysis was performed using SPSS version 26.0 software. Frequency with percentage was used to report categorical variables, while mean with standard deviation was used to report quantitative variables. Adjusted odds ratio with 95% confidence interval was used to identify factors associated with the outcome variable in the logistic regression model, considering  $P < 0.05$  to declare statistical significance.

### **RESULTS**

In this study, about 65% of nurses working in the dialysis unit had good practice. Nurses' years of experience [AOR: 11; 95% CI (1.29-39.3)], having training [AOR: 8.5; (95% CI; 2.26-27.5)], knowledge [AOR: 12 95% CI: 2.65-56.2)] were significantly associated with the practice of nurses regarding hemodialysis care.

## **CONCLUSION**

In this study we found that the nurse's practice towards hemodialysis service was good. Training, experience and knowledge of nurses were significantly associated with the experience towards hemodialysis care. Thus, interventions should focus on improving experience, awareness, and training of nurses to improve the practice of nurses regarding hemodialysis care.

## **KEYWORDS**

Hemodialysis; Nurses; Practice; Associated factors

## **ABBREVIATIONS**

AOR: Adjusted Odds Ratio

HD: Hemodialysis

RRT: Renal Replacement Therapy

AV: Arteriovenous

CI: Confidence Interval

USA: United States of America

## **INTRODUCTION**

Hemodialysis (HD) is an extracorporeal therapy that is prescribed to reduce the signs and symptoms of uremia and to substitute the partial functions of the kidneys [1]. It is the process of cleansing the blood from accumulated waste products. The process happens through the bidirectional movement of particles across a semipermeable membrane. Clinically, this movement takes place in and out of blood, across a semipermeable membrane. If the blood is exposed to an artificial membrane outside of the body, the process is HD [2].

HD has existed for more than 50 years and has prolonged the lives of millions of patients with renal failure worldwide. Although the fundamental principles of HD are still applying today, dialysis technology has technologically advanced markedly [3]. It is rapidly becoming more widespread as it has the potential to dramatically make better the quality of life and survival rate of chronic dialysis patients [4]. Regardless of many technical improvements in medical care and dialysis delivery that over the last few years, mortality and morbidity rates keep on high and the quality of life of uremic patient is still deprived and there are still important acute complications that had faced by nurses responsible for patients receiving HD in both acute and chronic clinical settings [5].

Generally, Complications of hemodialysis can be divided into two major groups: Treatment-related medical complications: Those are frequent intradialytic complications, these (in order of frequency) include hypotension, muscle cramps, nausea and vomiting, flushing of the face, headache, increased pruritus, chest pain, fever, and chills. The other is Machine-related complications: These complications are due to accidents or failure of safety mechanisms of hemodialysis treatment. They include air embolism, hemolysis, hyperthermia or hypothermia, blood loss, and conductivity problems [6]. The complications of hemodialysis are due to the disease itself as well as the mode of renal replacement therapy (RRT) [6]. Hemodialysis remained the most common treatment modality, with approximately 1,929,000 patients undergoing hemodialysis, for 89% of all dialysis patients. In

Egypt, the total prevalence of patients on dialysis is 264 per million [7]. The incidence and prevalence of end-stage organ failure in Ethiopia, including those of end-stage renal disease, are not known [8].

Quality has become an increasingly predominant part of our lives. The patient is constantly looking for quality care and services [9]. In nursing care of hemodialysis patients, quality is a complex concept that has multiple perspectives including the technical and personal aspects also; it should be conforming to standards of care, which are considered as a starting point for better practice. A patient undergoing hemodialysis procedure requires specially trained staff and special nursing care during phases of dialysis and the termination phase [10]. When appropriate dialysis care is available early in the course of management, the potential for better health improves, and that patient can remain active in family and community life [11]. Nurses are responsible for the direct care of patients undergoing dialysis. Dialysis nurses must have knowledge and skills because they reflected vital features of quality nursing care in the hemodialysis treatment [12]. Little is known about the nurses' practice and associated factors while caring for patients undergoing hemodialysis in Ethiopia. The current study was aimed to determine level of practice and associated factors of hemodialysis among nurses.

## **METHOD AND MATERIALS**

This study was conducted in Addis Ababa, the capital city of Ethiopia. The city has a subtropical highland climate. According to the 2017 estimation, Addis Ababa has a population of 6.6 million people. The city has 52 hospitals, of which 13 are public and 39 Private. Currently, there are 30 hemodialysis centers with a total of 186 hemodialysis chairs and approximately 800 patients on hemodialysis. Four of them are governmental dialysis units, with 31, 7, 5, and 3 hemodialysis chairs available in St. Paul hospital millennium medical college, Minillik II memorial hospital, Zewuditu memorial hospital, and Tikur Anbesa hospital respectively.

An Institution-based cross-sectional study was carried out on hemodialysis care provider nurses from all governmental hospitals dialysis units situated in Addis Ababa, Ethiopia from March 2020 to May 2020. A total of 62 nurses were included in the study.

A structured self-administered questionnaire was utilized to collect the data. Data were collected on different socio-demographic and other pertinent variables used to determine the practice of nurses regarding hemodialysis care. The mean score was utilized as a cut of point to describe the nurse's practice regarding hemodialysis care. A supervisor and two data collectors, who had a B.Sc. degree in nursing, were recruited to assist in the data collection process. Training was given for the supervisor and data collectors on the objectives of the study, the questions, and extent of explanations, the way to keep privacy and confidentiality and other ethical issues. The principal investigator was checked and reviewed the completeness of questionnaires and offered the necessary feedback for the supervisor and data collectors at spot to prevent bias. Data was checked for its completeness; then coded and entered into Epi Data 4.62 version. Finally, data was exported to SPSS version 25.0 for analysis. Categorical variables were presented using frequency with percentages, while quantitative variables were presented in medians with standard deviations. Bivariate regression analysis was done to identify candidate variables for the final model. Multivariate regression analysis was conducted to determine the association between dependent and independent variables. Adjusted odds ratio with 95% confidence interval was used to report factors associated with the outcome variable, considering  $P < 0.05$  to declare statistical significance.

## RESULTS

In this study, among 62 nurses working in different haemodialysis centres in Addis Ababa governmental hospitals, 60 nurses were participated with the response rate of 96.7%. Majority of participants (70%) were females, and the mean age of the respondents was 29.6±4.12 years. Regarding educational background of the participants, 93.3% of them were BSc holders followed by 5% master's degree (Table 1).

**Table 1:** Socio-demographic character of nurse's working in dialysis center of governmental hospitals in Addis Ababa 2020.

Variables	Category	Frequency	Precent
Sex	Male	18	30
	Female	42	70
Age	<30 years	39	65
	≥30 years	21	35
Marital status	Single	29	48.3
	Married	27	45
	Divorced	3	5
	Separated	1	1.7
Educational level	Diploma	1	1.7
	B.Sc.	56	93.3
	Masters	3	5
Work experience	<4 years	24	40
	4 years and above	36	60

In the current study, it was reported that 65% of nurses working in the dialysis room of governmental hospitals in Addis Ababa had good practice, followed by 35% of them having poor practice (Table 2).

**Table 2:** Knowledge of nurses working in dialysis center of governmental hospitals in Addis Ababa, 2020.

Variables	Category	Frequency	Perce nt
Practice of nurses	Good	39	65
	Poor	21	35
Were measure the weight of a patient to determine amount of fluid to be removed during dialysis	Yes	50	83.3
	No	10	16.7
Were take vital signs like BP for hemodynamic status, temperature for sepsis and respiration for fluid overload?	Yes	60	100
	No	0	0
Were Ask the patient whether take or not antihypertensive medication, especially if systolic blood pressure is below 100 mmHg	Yes	43	71.7
	No	17	28.3
Were Asses vascular access site for infection	Yes	46	76.7
	No	14	23.3
Were they Sanitizing the machines before starting dialysis?	Yes	60	100
	No	0	0
Was hand hygiene performed before putting on gloves?	Yes	30	50
	No	30	50
Were clean gloves put on before removing the old CVC dressing?	Yes	60	100
	No	0	0
Were gown, protective eyewear and mask or face shield worn during CVC exit site care or before annulation?	Yes	25	41.7
	No	35	58.3
Was a new sterile dressing applied to the CVC exit site?	Yes	54	90
	No	6	10
CVC exit site cleaned with antiseptic beginning at the exit site and working outward toward the periphery?	Yes	59	98.3
	No	1	1.7
Was the Chlorhexidine or antiseptic allowed to dry before applying the clean dressing?	Yes	60	100
	No	0	0
Were the CVC connections disinfected by scrubbing the hub thread for 15 seconds with an alcohol pad and allowed to dry?	Yes	57	95
	No	3	5
Were sterile syringes connected to each port with an aseptic technique?	Yes	57	95
	No	3	5

Was each port aspirated the heparin and check the access for patency?	Yes	60	100
	No	0	0
Were annulations needles inserted aseptically?	Yes	57	95
	No	3	5
Was treatment initiated with aseptic connection to blood lines	Yes	59	98.3
	No	1	1.7
Was the treatment initiated successfully?	Yes	60	100
	No	0	0
Was hand hygiene performed after the CVC initiation or annulation procedure was complete?	Yes	59	98.3
	No	1	1.7
Was hand hygiene performed before putting on gloves?	Yes	46	76.7
	No	14	23.3
Were gown, protective eyewear and mask or face shield worn during the CVC or AVF discontinuation procedure?	Yes	29	48.3
	No	31	51.7
Was reduce the pump speed up to 100-150 ml/mints and stop it?	Yes	60	100
	No	0	0
Were clean gloves put on before starting the dialysis discontinuation?	Yes	59	98.3
	No	1	1.7
Was the extracorporeal circuit rein fused?	Yes	60	100
	No	0	0
Were the bloodlines properly disconnected?	Yes	60	100
	No	0	0
Were sterile port caps applied post treatment using aseptic technique and apply dressing?	Yes	60	100
	No	0	0
Were needles removed with aseptic technique?	Yes	60	100
	No	0	0
Were needles discarded in the sharps container?	Yes	60	100
	No	0	0
Were needle sites held with clean gauze using clean-gloved hands?	Yes	60	100
	No	0	0

In this study, 60% of nurses working in dialysis unit of hospitals had taken training regarding dialysis procedure and intradialytic complication management, and 66.6% of the nurses who had taken the training reported that they were satisfied by the training (Table 3).

**Table 3:** Factors associated with practice of nurses working in dialysis center of governmental hospitals of Addis Ababa, 2020.

Factors	Categ ory	Freque ncy	Perce nt
Have you received training related to dialysis procedure and intradialytic complication management during your professional development?	Yes	36	60
	No	24	40
If yes, are you satisfied with the training?	Yes	24	66.6
	No	12	33.3
Do you have a dialysis procedure and intradialytic complication management guideline or standard in your organization?	Yes	32	53.3
	No	28	46.7
If yes Q (403), are you read the guidelines.	Yes	28	87.5
	No	4	12.5
Do you have a dialysis Patient assessment form in your organization?	Yes	38	63.3
	No	22	36.7
Do you have a form for assessment of A-V fistula or central venous catheter site in your organization?	Yes	37	61.7
	No	23	38.3
Do you have fluid assessment form in your hospital?	Yes	15	25
	No	45	75
Is their assigned dietician in your hospital?	Yes	0	0
	No	60	100

### ***Practice related Factors of Nurses towards Hemodialysis***

In the bivariable logistic regression analysis, experience of nurses, training of nurses regarding hemodialysis, knowledge of nurses, presence of patient assessment guideline, presence of AV-fistula assessment form was associated with the practice of nurses. Those variables that have a p-value less than or equal to 0.25 were entered

to a multivariable logistic regression model to adjust for possible confounders. In the multivariable logistic regression analysis, experience of nurses, training of nurses and knowledge of nurses regarding hemodialysis were found to be significantly associated with the practice of nurses. Nurses who had 4 years and above experience in working dialysis unit were 11 times more likely to have good practice compared to those who had less than 4 years' experience [AOR: 11; 95% CI (1.29-39.3)]. Nurses who have been taken training regarding hemodialysis were 8.5 times more likely to have good practice compared to those who don't have been taken training regarding hemodialysis [AOR: 8.5; (95% CI; 2.26-27.5)]. In this study, nurses who were knowledgeable regarding hemodialysis were 12 times more likely to have good practice compared to those nurses who were not knowledgeable [AOR: 12(95% CI; 2.65-56.2)].

**Table 4:** Bivariable and multivariable logistic regression on practice of nurses working in dialysis center of governmental hospitals in Addis Ababa, 2020.

Variables		Knowledge		COR (95%)	AOR (95%)
		Good	Poor		
Experience of nurses	< 4 years	6(10)	18(30)		1
	4 years and above	33(55)	3(5)	13.3(2.85-49.9)	11(1.29-39.3)*
Training of nurses	Yes	33(55)	3(5)	11(2.99-41.8)	5(1.12-22.2)*
	No	6(10)	18(30)		1
Knowledge of nurses	Good	37(61.7)	5(8.3)	15.9(10.3-73.3)	12(2.65-56.2)*
	Poor	2(3.3)	16(26.7)		1
Presence of patient assessment guideline	Yes	34(56.7)	4(6.7)	28(6.86-121)	8.6(0.6-24.12)
	No	5(8.3)	17(28.3)		1
Presence of fluid assessment guidelines	Yes	14(23.30)	1(1.7)	11(1.35-29.5)	0.34(0.67-22.3)
	No	25(41.7)	20(33.3)		1
Presence of AV - fistula assessment form	Yes	32(53.3)	5(8.3)	14.6(4.00-53.41)	1.32(0.23-21.1)
	No	7(11.7)	16(26.7)		1

## **DISCUSSION**

This facility based cross sectional study was conducted to assess practice and associated factors regarding hemodialysis care among nurses. Among the participants, the majority (70%) were females, which shows the dominance of females over the males. The mean age of the respondents was  $29.6 \pm 4.12$  years; shows that the newly graduate and expert nurses were interested to the hemodialysis care practice. These findings were similar to the study done in USA, found that the most respondents in their study were young age [13].

The majority (93.3%) of participants had bachelor's degree and 5% had masters and only 1.7% diploma; this is due to the fact that people are upgrading themselves and majority of the participants were from teaching and tertiary hospitals that were administered by university, since the university criteria to be recruited is bachelor degree holders.

In this study, it was reported that 65% of nurses working in the dialysis care had good practice, followed by 35% of nurses had poor practice. A study conducted in Saudi Arabia reported that 84.13% of participants had good practice, which is higher than our finding [13]. Similarly, results from a study conducted in Khartoum, Sudan reported that 78% of nurses had good practice. However, a study finding in Brazil reported that 47.54% of nurses had good practice which is by far lower than our finding [5]. These differences may be due to variation in

knowledge, training and experience among nurse and the availability of facilities and services in those various set ups.

In this study, nurses who had 4 years and above experience in working dialysis unit were 11 times more likely to have good practice compared to those who had less than 4 years' experience. This finding is supported with the results of study conducted by Hosien Shahdadi and Mozghan Rahnama that the nurse's experience had positive effect on the outcome of patients within hemodialysis [14], and other study conducted in Iran supports the this finding [15]. This is because experience improves the confidence, protection, and to gain patients' trust and ensure a successful dialysis through adaptation of practice of care.

In the current study, we found that nurses who took training regarding hemodialysis were 8.5 times more likely to have good practice compared to those who did not took training regarding hemodialysis care. This finding was agrees with the results of studies done in Iran [15] and Saudi Arabia [16]. This is because of provision of specialized training for nurses improve their skills to manage complication and prevent infections.

In this study, nurses who were knowledgeable regarding hemodialysis were 12 times more likely to have good practice compared to those nurses who were not knowledgeable. This finding was in line with the results of studies conducted in Iran (Iran), Saudi Arabia [16] and Jordan [17]. The possible justification could be as nurses gain basic knowledge on hemodialysis to provide high quality care, may take advantage of their technical knowledge.

### **LIMITATION OF THE STUDY**

The current study could not establish cause-effect relationship given that it was a cross-sectional study. Use of small sample size was also another limitation of the study that may affect the generalizability of the study finding.

### **CONCLUSION**

In the current study the practice of nurses towards hemodialysis care was good. The year of experience, having getting training and their level of knowledge had shown significant association with practice among nurses on hemodialysis care. Interventions should focus on providing training and recruiting experienced nurses with good level of knowledge on hemodialysis care to maintain and improve patient's health status.

### **ACKNOWLEDGEMENTS**

We are indebted to Addis Ababa University, College of Health Sciences for its financial support for data collection. We are also thankful for the hospitals involved in this study for their cooperation. Finally, the authors are deeply thankful to the supervisors and data collectors.

### **AUTHORS' CONTRIBUTIONS**

TB, AF, conceived and designed the study and developed the data collection instruments. TL and TL performed the statistical analysis and wrote all versions of the manuscript. All authors critically revised and approved the final manuscript.

### **COMPETING INTERESTS**

The authors declare that they have no competing interests.

## **REFERENCES**

1. Gilbert SJ, Weiner DE, Gipson DS et al. (2017) National kidney foundation's primer on kidney diseases sixth edition. Bone Disorders in Chronic Kidney Disease 276-287.
2. Kallenbach JZ (2019) Review of hemodialysis for nurses and dialysis personnel - E-book.
3. Oliver J (2013) Comprehensive clinical nephrology. Journal of Chemical Information and Modeling 53(9).
4. Lok CE and Mokrzycki MH (2011) Prevention and management of catheter-related infection in hemodialysis patients. Kidney International 79(6): 587-598.
5. Ashby D, Borman N, Burton J et al. (2019) Renal association clinical practice guideline on haemodialysis. BMC Nephrology 20: 1-36.
6. Ahmad S (2009) Clinical manual dialysis. Springer Science & Business Media.
7. Al-Mawsheki E, Ibrahim MH, Taha NM (2016) Nurses' knowledge and practice regarding care for the patients during hemodialysis. The Medical Journal of Cairo University 84(1): 1135-1141.
8. Ahmed MM, Tedla FM, Leichtman AB et al. (2019) Organ transplantation in Ethiopia. Transplantation 103(3): 449-451.
9. Mosadeghrad AM (2012) A conceptual framework for quality of care. Materia Socio-Medica 24(4): 251.
10. Saleh M, Ali J, Afifi W (2018) Nurses compliance to standards of nursing care for hemodialysis patients: Educational and training intervention. IOSR Journal of Nursing and Health Science (IOSR-JNHS) 7(2): 48-60.
11. El-Moghazy G, Hassan ASA, Sorour A et al. (2013) Nurses knowledge and practice regarding intradialytic complications for hemodialysis patient. Journal of American Science 9(11): 300-307.
12. Price CA and Paganini EP (1999) Debate: should nurses have a larger role in the outpatient dialysis setting than they currently do?. In Seminars in Dialysis: ADVANCES Meeting Report 12(5): 359-362.
13. Alramadhan ES, Alsayed S, Alshalawi AA et al. (2019) Assessment of the nurses' skills during caring for hemodialysis patients. Austin Journal of Nursing & Health Care 6(1): 1-5.
14. Shahdadi H and Rahnema M (2018) Experience of nurses in hemodialysis care: A phenomenological study. Journal of Clinical Medicine 7(2): 30.
15. Manandhar DN, Chhetri PK, Poudel P et al. (2017) Knowledge and practice of hemodialysis amongst dialysis nurses. Journal of Nepal Medical Association 56(207): 346-351.
16. Al Qahtani AS and Almetrek MA (2017) Knowledge, attitudes and practice of nurses in renal dialysis units regarding infection control in Abha city-Saudi Arabia, 2015. The Egyptian Journal of Hospital Medicine 66(1): 103-114.
17. Alnawafleh KA, Mohammad ZA, Al-Nwafleh HA (2018) Auditing of Nursing care for hemodialysis patients at Ma, an Governmental hospitals-Jordan. Assiut Scientific Nursing Journal 6(15): 89-98.