

CLINICAL REVIEW

---

# What is the Impact of COVID-19 Pandemic on the RCH (Reproductive and Child Health) Programme in Rajasthan, Because of Nationwide Lockdown (April 2020 to June 2020)?

Piyush Kumar<sup>1\*</sup> and Habib Hasan Farooqui<sup>2</sup>

<sup>1</sup>*Health Department, Government of Bihar, Bihar, India*

<sup>2</sup>*Public Health Department, Indian Institute of Public Health, Delhi, India*

Correspondence should be addressed to Piyush Kumar, Health Department, Government of Bihar, Bihar, India

Received: 11 March 2022; Accepted: 16 April 2022; Published: 24 April 2022

Copyright © Piyush Kumar. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## **ABSTRACT**

### **BACKGROUND**

The proper, timely, adequate delivery of effective and high-quality child health and reproductive (RCH) services is of greatest significance and utmost priority mainly because of situations erupting from the current ongoing pandemic of COVID-19 as well as other cofactors in the state of Rajasthan, India. This significance and priority are particularly due to many factors such as huge as well as increasing population with limited qualified, skilled human resources.

### **OBJECTIVE**

The key objective is to find out the effects of SARS-CoV-2/COVID-19 pandemic on the Reproductive and Child Health Programme of Rajasthan in India over the 3 months after lock down enforcement nationwide in March 2020.

### **SETTINGS/DESIGN**

Different indicators of RCH programme i.e., immunisation, maternal and child health, family planning) for Rajasthan were observed, studied, collected and compared with previous year 2019 for the period of April/May/June -2019 and 2020. The percent increase and downfall is observed, calculated, presented, from the available data to find out the status of delivery of essential RCH health services. The need assessed and percent of achievement of assessed need is also compared, and future achievement projected as per achievement.

## **MATERIALS AND METHOD**

To know the effect of pandemic era on RCH health programme of Rajasthan during the period from 3 months after lock-down announced in the month of March 2020 i.e. April/May/June 2020 in Rajasthan the data observed, calculated and obtained from HMIS (Health Management Information System- of Ministry of Health and Family Welfare (MoHFW), Government of India for RC Health programme, (Reproductive and Child Health).

## **RESULT**

The results of data analysis for RCH services, functioning in the state of Rajasthan during the month during the period from April to June 2020 (during the period from lockdown) as compared to 2019 (no lockdown, for same duration suggest that the immunisation services were badly affected during the period from the lockdown period in 2020. It is possible that newborn children and the older ones have not received/delivered proper immunisation services during the period from the lockdown as evident from the data observation-analysis.

## **CONCLUSION**

India and different states have launched several plans and strategies to deal with COVID-19 pandemic. However, this study report found that insufficient attention has been given to very essential services of RCH in Rajasthan as evident from the results of this study. The problems in the delivery of healthcare services should be dealt with properly, timely, adequately added with a proper update of the latest dynamic plan to carry on essential health delivery services even in pandemics and other situations of emergencies. Rajasthan should develop an exclusive updated latest dynamic plan to deal with situations to guarantee delivery of very essential services such as RCH during the period from pandemics or any other natural calamities-emergencies.

The ongoing COVID-19 pandemic has disclosed the vulnerability of women and children and at the same time teaches us about the importance of equity in healthcare. The children and women who are considered most vulnerable in emergencies and distress must have special supportive healthcare all the time especially during the period from pandemics-emergencies and other natural disasters.

## **KEYWORD**

Health; Decrease; Pandemic; Services/service; COVID-19; Sars-Cov-2

## **ABBREVIATION**

RCH: Reproductive and Child Health; HCWs: Health Care Workers; ANC: Antenatal Care; BMJ: British Medical Journal; VPD: Vaccine Preventable Diseases; NFHS: National Family Health Survey; WHO: World Health Organization; TFR: Total Fertility Rate; UNICEF: United Nations International Children's Emergency Fund; ANM: Auxiliary Nurse Midwife; LHV: Lady Health Visitor; MCP: Mother and Child Protection; LMICs: Low-Income and Middle-Income Countries; LiST: Lives Saved Tool; HMIS: Health Management Information System; MoHFW: Ministry of Health and Family Welfare; BCG: Bacillus Calmette Guerin; DPT: Diphtheria Pertussis Tetanus; DT: Diphtheria Tetanus; MR: Measles Rubella; TT: Tetanus Toxoid; IUD: Intrauterine Device; OPV: Oral Polio Vaccine

## **INTRODUCTION**

The proper, timely, adequate delivery of effective and high-quality RCH (Reproductive and Child Health) services is of greatest significance and utmost priority in situations erupting from the current ongoing pandemic of COVID-19 as well as other cofactors in the state of Rajasthan, India. This significance and priority are particularly due to many factors such as huge as well as increasing population with limited qualified, skilled human resources [1], vast area, difficult geographical locations, underdeveloped limited infrastructure as well as huge demand on healthcare system due to growing burden of NCD (non-communicable) and CD (communicable diseases). The women and children are considered as vulnerable groups affected by hazards of such disaster conditions like COVID-19 pandemic; hence this topic needs essential and special attention especially in current pandemic scenario 2020-2021. India is second most populous nation after China and women's with children's constitutes a large section of current population [2].

The current ongoing COVID-19/SARS-CoV-2 pandemic situation has presented a big challenge even for highly developed healthcare systems around the world. A sense of fear and insecurity has gripped the whole world's populations since 2019 due to the pandemic and the state of Rajasthan in India is not an exception. The limited underdeveloped healthcare system and poor health and other resources including transportation (ambulance services) manpower, infrastructure, etc. have been largely utilized to deal with the situation erupting from the SARS-CoV2. This one-sided shifting of resources added with several other factors has incredible-sensational consequences on progress of various health programmes such as RCH (Reproductive and child health) running previously before this COVID-19 pandemic era. Historically also the children and women usually suffer more in such vulnerable situations like disasters and pandemics. One such finding is given by study done by Alice Reid on effects of 1918-1919 influenza pandemic he found that young children and Infants are a more vulnerable group to different forms of communicable disease, and their relative mortality was also increased abnormally in pandemic but talk about the young is mostly limited to mentions of school closures, or in the background of illness or mortality to whole families [3].

Children and pregnant mothers are vulnerable groups due to their special health requirements and need more attention in disasters and pandemics due to special health and safety requirements such as ANC (antenatal care) and immunisation. After births newborn are naturally immune to several diseases because they have antibodies acquired from their mothers in a natural way. However, this acquired immunity declines slowly and gradually with time as the baby grows up. Hence immunisation/vaccination is required to protect them from various vaccine preventable diseases (VPD) which are provided through universal immunization programmes (UIP-1985) to all children in India (Universal Immunization Programme (UIP) (Table 1) [4].

The encouragement for child and maternal health, women upliftment has been done and addressed by many organisations of India and international repute for several years mentioning them as special vulnerable groups in times of disaster such as pandemic. The SARS-CoV-2 pandemic seems to disclose the limitations of the health system to save the above-mentioned special groups as reported by WHO and other agencies expressing concerns over several issues related to women and children's [5]. As per current pandemic situation it is of special focus and requirement to have a separate health cadre of trained personnel in the field of RCH to protect child and mother health hazards in special situations of disasters like COVID-19 pandemics. It seems many protective and

advantageous essential health services are not being provided in COVID-19 era which may produce undesirable and detrimental effects at mass level in the coming future. The World Health Organisation (WHO) have addressed these and many other concerns such as COVID-19 causes disruptions to child protection services in more than 100 countries, UNICEF (United Nations International Children's Emergency Fund) survey, millions more cases of female genital mutilation, unintended pregnancy, violence, child marriage, expected due to the COVID-19 pandemic [6].

<b>National Immunization Schedule</b>				
<b>Vaccine</b>	<b>When to give</b>	<b>Dose</b>	<b>Route</b>	<b>Site</b>
<b>For Infants</b>				
BCG	At birth or as early as possible till one year of age	0.1ml (0.05ml until 1 month of age)	Intra -dermal	Left Upper Arm
Hepatitis B Birth dose	At birth or as early as possible within 24 hours	0.5 ml	Intramuscular	Anterolateral side of mid-thigh-LEFT
OPV Birth dose	At birth or as early as possible within the first 15 days	2 drops	Oral	-
OPV 1,2 and 3	At 6 weeks, 10 weeks, and 14 weeks	2 drops	Oral	-
IPV (Inactivated Polio Vaccine)	14 weeks	0.5 ml	Intramuscular	Anterolateral side of mid-thigh-RIGHT
Pentavalant 1,2 and 3	At 6 weeks, 10 weeks, and 14 weeks	0.5 ml	Intramuscular	Anterolateral side of mid-thigh-LEFT
Rota Virus Vaccine	At 6 weeks, 10 weeks, and 14 weeks	5 drops	Oral	-
Measles 1 <sup>st</sup> Dose	9 completed months-12 months. (Give up to 5 years if not received at 9-12 months age)	0.5 ml	Subcutaneous	Right Upper Arm
Vitamin A, 1 <sup>st</sup> Dose	At 9 months with measles	1 ml (1 lakh IU)	Oral	-
<b>For Children</b>				
DPT 1 <sup>st</sup> Booster	16 months - 24 months	0.5 ml	Intramuscular	Anterolateral side of mid-thigh-LEFT
OPV Booster	16 months - 24 months	2 drops	Oral	-
Measles 2 <sup>nd</sup> Dose	16 months - 24 Months	0.5 ml	Subcutaneous	Right Upper Arm
Vitamin A (2 <sup>nd</sup> to 9 <sup>th</sup> Dose)	16 months with DPT/OPV booster, then, one dose every 6 month up to the age of 5 years)	2 ml (2 lakh IU)	Oral	-
DPT 2 <sup>nd</sup> Booster	5-6 years	0.5 ml	Intramuscular	Left Upper Arm
TT	10 years and 16 years	0.5 ml	Intramuscular	Upper Arm

**Table 1:** Vaccination schedule under the UIP.

To recognize the real effect of COVID-19 on child health, maternal, and family planning services provided under RCH programme the service utilisation data and other outcome indicators were analysed during month of April to June 2020 (lockdown phase) with non-lockdown phase of during the period from the same months in the previous year 2019. India was also under a lockdown period from 25<sup>th</sup> March 2020 to 31<sup>st</sup> May 2020 for a total period of 68 days during the period from which public transportation and many other services were stopped. The Ministry of Home Affairs, Government of India had issued order number 40-3/2020-D dated 24/03/20 regarding guidelines on suppression of COVID-19 epidemic in the country stating in point number 6 that all transport services i.e. air, rail, roadways will remain suspended with few exceptions [7]. The HCWs (health care workers) not having their own vehicles to do routine duties added with high rate of COVID-19 infection and deaths of HCWs also seems to affect healthcare delivery services.

## **LITERATURE REVIEW**

RCH (Reproductive and child health) programme is a matter of concern for India even before independence due to several factors like socio-economic status, inequities, and lack of resources, infrastructure and many more. The practical problems in delivering these services are also faced at all levels. The RCH services are a matter of greatest concern in the ongoing pandemic era since December 2019 at global levels and particularly for countries like India [8]. These concerns have been taken seriously by international organisations like WHO/UNICEF and alert regarding immunisation, ANC is being delivered at various levels as well as much scientific research is being carried out in this regard. In one such study as informed by WHO that Pregnancy with SARS-CoV-2 is not as much likely than non-pregnant women with SARS-CoV-2 to have symptoms, but more likely to need special-intensive care if severely ill, as per most recent research findings of the BMJ (British Medical Journal) on the risks of SARS-CoV-2 for babies and pregnant women. The text literature suggests that pregnant women seen at the hospital with supposed or confirmed SARS-CoV-2 are less likely to encounter a fever or muscle pain, but if they develop severe illness, they are more likely to need ICU treatment than non-pregnant women with SARS-CoV-2 [9]. The family planning programme is equally important as the population of India is increasing at a very fast pace which is increasing more demand on available resources. The key purpose of this review is to find out if there is convincing evidence that Rajasthan RCH programme is getting affected by SARS-CoV-2 pandemic. In the same statement mentioned before WHO, also stated that it is important to recognise the increased anxiety and stress caused by SARS-CoV-2 which may be predominantly felt by children, pregnant women, their partners, in recent times-pregnant women, and families. Therefore, healthcare workers have a big role in providing care to pregnant women in an appropriate and benevolent way.

The scope of this review is limited as more data and materials are required to complete it and there is also a word limit for this report. As per NFHS-4 (National Family Health Survey) survey data the total fertility rate (TFR) in Rajasthan is 2.4 children per woman, which is more than replacement level fertility (2.1). Fertility declined by only 0.4 children in the last 13 years between NFHS-1 and NFHS-3 and has decreased further by 0.8 children in the 10 years period between NFHS-3 and NFHS-4.

Regarding pregnancy outcomes the NFHS-4 finding were that 91 percent of previous pregnancies in the last five years preceding the survey resulted in a live birth, and the rest 9 percent ended in foetal wastage (miscarriage, or stillbirth, abortion) and among this miscarriage was the most commonly found responsible for foetal wastage, about 7 percent of all pregnancies, and abortions lead to 2 percent foetal wastage. NFHS-4 survey also found that most of the abortions were done in the private health sector (59 percent) and only 21 percent were done in the public health sector. Added to this report of NFHS-4 is that one-fourth of women having an abortion, reported to experience complications from the abortion. NFHS-4 survey found that awareness of contraception is almost universal in Rajasthan, except some methods are still less popular. 81 percent of recently married women know about injectable methods of contraception while 11 percent know about female condoms. Among all women, only 45 percent have knowledge of emergency contraception.

Regarding antenatal-care the NFHS-4 survey reported that among pregnant women's who gave birth in the last five years before the survey, more than 4/5<sup>th</sup> (83 percent) received ANC for their last birth from a health professional (28percent from an auxiliary nurse midwife, and lady health visitor (LHV), nurse, or midwife 55

percent from a doctor). 14 percent did not get any ANC. Among mothers who gave birth in the last 5 years before the survey, more than 4/5<sup>th</sup> (85 percent) registered the pregnancy for the most recent live birth. Among the registered pregnancies, 92 percent received an MCP Card (Mother and Child Protection Card). Only 39 percent of mothers in Rajasthan acknowledged at least 4 ANC visits for their last birth. Furthermore, regarding delivery care the survey found that more than 4/5<sup>th</sup> of births (84 percent) take place in a health facility (mostly a government facility) and 16 percent at home. The survey found that the percent of births in a health facility tripled in the 10 years period between NFHS-3 and NFHS-4, from 30 percent in NFHS-3 to 84 percent in NFHS-4. Institutional births are more common among women who have received an ANC check, women who are having their first birth, women with 12 or more years of schooling, urban women, and Sikh women. Survey also found that 84 percent of children in Rajasthan who were born in the last five years were born in a health facility [10].

Regarding immunization/vaccination the survey reports that more than half (55 percent) of children between 12 months - 23 months get all available vaccinations against 6 (tuberculosis, diphtheria, pertussis, tetanus, polio, and measles,) important childhood diseases before the survey. Most of the children are at least partially vaccinated; only 7 percent have not got any vaccinations at all. The survey also found that 89 percent of children have got a BCG vaccination. Added to this, very less children have received other basic vaccinations (65 percent have got at least the recommended three doses of polio vaccine, 72 percent have got the three recommended doses of DPT vaccine, and 78 percent have been vaccinated against measles). There is a remarkable dropout between the 1<sup>st</sup> and 3<sup>rd</sup> doses of DPT vaccine (from 86 percent to 72 percent) and polio vaccine (from 88 percent to 65 percent). One remarkable finding is that Coverage with all basic vaccinations is high for children whose mothers have completed 12 or more years of schooling.

## **RATIONALE**

One study published in the Lancet Global Health [11] by Timothy Robertson et al. about the indirect effects of SARS-CoV-2 on maternal and child mortality in LMICs with use of the Lives Saved Tool (LiST), found that reduced provision and use of reproductive, maternal, newborn, and child health-care services might substantially increase maternal and child deaths but this dissertation project is a totally different study incorporating three aspects viz. Maternal health, child health and family planning based on real-time data from an accredited source to find out the impact of SARS-CoV-2 on RCH through output health indicators. This study, furthermore, addresses the situations of Rajasthan, the constraints and limited access to information means that some major figures, such as for India and several other states and union territories as well as for longer periods data is not available-not been included [12-16]. Finally, although several articles on this subject are available as seen in Google search, to evaluate the impact of SARS-CoV-2 on mother, child etc through studies using either qualitative or quantitative methods, to measure the effectiveness of RCH programs during the period from SARS-CoV-2 is comparatively scarce. I have included relevant examples of these studies being done by myself for some other states and India [17-21], yet it can be said at the outset that the dearth of such studies needs to be redressed by the research community. Previous studies mentioned above have found a negative impact on health indicators of RCH programmes for few states and India which aroused my interest to know the impact of SARS-CoV-2 on other states. Rajasthan is the biggest state in India, so I decided to study Rajasthan to know how RCH programmes perform in such a big state during the lockdown period from the above-mentioned period.

## **AIM**

RCH (Reproductive and child health) programme is a matter of concern for India even before independence due to several factors like socio-economic status, inequities, and lack of resources, infrastructure and many more. The key aim of this study is to understand the effect of COVID-19 pandemic era on RCH programme of Rajasthan for 3 months after lock down declared in the month of March 2020 i.e. April/May/June 2020 in Rajasthan.

## **OBJECTIVE**

To know the real effect of SARS-CoV-2 on child health, maternal health, and family planning services provided under RCH programme the service utilisation data and other outcome indicators were analysed during the month of April to June 2020 (lockdown phase) with non-lockdown phase of during the period from the same months in the previous year 2019. India was also under a lockdown period from 25<sup>th</sup> March 2020 to 31<sup>st</sup> May 2020 for a total period of 68 days during the period from which public transportation and many other services were stopped. The Ministry of Home Affairs, Government of India had issued order number 40-3/2020-D dated 24/03/20 regarding guidelines on containment of SARS-CoV-2 epidemic in the country stating in point number 6 That all transport services i.e., air, rail, roadways will remain suspended with few exceptions.

## **METHODS**

### ***Study Design***

The study design is retrospective mixed quantitative and qualitative analysis of the RCH programme of Rajasthan.

### ***Study Setting and Duration***

To know the impact of pandemic era on RCH programme of Rajasthan during 3 months after lock down declaration in the month of March 2020 i.e., April/May/June 2020 in Rajasthan. The data period under consideration is for the year 2019 and 2020, which was analysed to report several outcome indicators of interest.

### ***Sampling Method***

The data obtained from HMIS (Health Management Information System) of Ministry of Health and Family Welfare (MoHFW), Government of India for RCH programmes (Reproductive and Child Health). The data analysis was done by Microsoft office software. The data is available for free distributions as mentioned by MoHFW-see excel table 2 below [12]. Different outcome indicators were taken into account to know the effect of SARS-CoV-2 Pandemic on delivery of health services to mother and child as well as family planning health services under the umbrella of RCH programme.

### ***Study Participants***

This study has not involved any human or animals in real or for experiments.

### ***Inclusion criteria***

Different indicators group of RCH programme (immunisation, maternal and child health, family planning) for Rajasthan during the month of April/May/June -2019 and 2020 (Table 2).

### Exclusion criteria

Different indicators group of RCH programme (immunisation, maternal and child health, family planning) for Rajasthan other than the month of April/May/June -2019 and 2020.

<b>Immunisation</b>
BCG-BACILLUS CALMETTE GUERIN
PENTA 3+DPT 3
DT (2 <sup>ND</sup> DOSE) OR DPT-5
MEASLES + MR
POLIO (OPV 3)
TETANUS TOXOID (TT10)
TETANUS TOXOID (TT16)
VITAMIN A (1 <sup>ST</sup> DOSE)
VITAMIN A (5 <sup>TH</sup> DOSE)
VITAMIN A (9 <sup>TH</sup> DOSE)

**Table 2:** Immunisation items Rajasthan.

### Study Tools and Measurements

The data analysis was done by Microsoft office software. The data is available for free distributions as mentioned by MoHFW (table 3) [12]. Different outcome indicators were taken into account to know the effect of SARS-CoV-2 Pandemic on provision of health services to mother and child as well as family planning health services under the umbrella of RCH programme.

	A. 5	9.1.12	Polio (OPV3)	1822630	Children upto 1 year of age	39,68,965
	A. 6	9.5.3	Tetanus Toxoid (TT10)	1615140	Children aged less than 10 years of age	15,99,129
	A. 7	9.5.4	Tetanus Toxoid (TT16)	1863310	Children aged less than 16 years of age	14,87,006
	A. 8	9.8.1	Vitamin A (1 <sup>st</sup> Dose)	1822630	Children upto 1 year of age	35,85,146
	A. 9	9.8.2	Vitamin A (5 <sup>th</sup> Dose)	1780150	Children 3 year of age	15,72,917
	A. 10	9.8.3	Vitamin A (9 <sup>th</sup> Dose)	1725000	Children 5 years of age	14,72,555
B. Maternal and Child Health	B. 1	1.1	ANC	2066780	Pregnant women	64,71,349
	B. 2	2.1.1a, 2.1.1b	Maternal Health - Home Deliveries			2,22,369
	B. 3	2.2	Maternal Health - Institutional Deliveries	1892660	Total Births	36,45,173
C. Family Planning	C. 1	8.13	Condom-User*	8995120	Unsterilised Eligible Couples	33,62,225
	C. 2	8.12, 8.14	Oral Pill Users*	8995120	Unsterilised Eligible Couples	26,10,843
	C. 3	8.3, 8.4, 8.5	IUD Insertions	8995120	Unsterilised Eligible Couples	9,20,990
	C. 4	8.2.1, 8.2.2, 8.2.3, 8.2.4, 8.1.1	Sterilisation	15220170	Total Eligible Couples	1,99,900
	C. 5	8.2.1, 8.2.2, 8.2.3, 8.2.4	Tubectomy	15220170	Total Eligible Couples	1,99,041
	C. 6	8.1.1	Vasectomy	15220170	Total Eligible Couples	859

**Table 3:** Source HMIS-MoHFW.

### ***Outcome***

The study outcome is presented in the form of tables, graphs and literature.

### ***Exposure***

All the relevance to the study is listed in reference as well as data source is also documented.

### ***Data Collection Tool***

The data obtained from HMIS (Table 3) of Ministry of Health and Family Welfare (MoHFW), Government of India for RCH programmes (Reproductive and Child Health) with the help of internet. The data analysis was done by Microsoft office software. For understanding impact on immunisation, the output indicators considered were immunisation data related to: (1) Tetanus Toxoid (TT16), (2) Vitamin A (1<sup>st</sup> Dose), (3) Vitamin A (5<sup>th</sup> Dose), (4) Vitamin A (9<sup>th</sup> Dose), (5) BCG (Bacillus Calmette Guerin), (6) Penta 3+DPT 3, (7) DT (2<sup>nd</sup> Dose) Or DPT-5, (8) Measles + MR, (9) Polio (OPV 3), (10) Tetanus Toxoid (TT10). For maternal health and family planning outcome indicators considered were: (1) ANC, (2) Home Deliveries, (3) Institutional Deliveries, (4) Condom User, (5) Oral Pill User, (6) IUD Insertion, (7) Sterilisation, (8) Tubectomy, (9) Vasectomy.

### ***Study Size, Place and Population***

Rajasthan is the biggest state of India with a population of 06.86 crores, as per census 2011, density 200/square kilometre, with some difficult geographic locations. The total numbers of districts are 33 with division into sub districts. The total child population as per 2011 census was (0 years - 6 years) 106.5 lakh and the total female population was 329.97lakh whereas males were 355.5 lakh in number. The population growth as per 2011 census was 21.31percent and state have 5.66 percent of total population of Rajasthan with literacy rate of 66.11percent. The projected population is 08.07 crores. The total married women as per 2011 census were 187.51 lakh with number of women with 7 plus child born was 15.57 lakh. Rajasthan is among the top 10 most populous states as well as growth rate [8].

### ***Data Collection Process (Settings and Design)***

Different indicators group (immunisation, maternal and child health, family planning) of RCH programme for Rajasthan were taken into consideration for the study and data collected and compared from previous year during the month of April/May/June -2019 and 2020.

### ***Data Analysis***

The percent increase and decline is calculated from the collected and available data to know the status of provision of important and essential health services. The need assessed and percent of achievement of assessed need is also compared, and future achievement projected as per achievement. The data analysis was done by Microsoft office software. The data is also shown in tabulated as well as graphical form for ease of understanding. All the data obtained were analysed using Microsoft office software.

### ***Ethical Consideration***

The ethical committee have been requested to provide clearance over the ethical issue as this study has not involved any human or animals in real or for experiments.

## **RESULTS**

Antenatal care is very much essential for safeguarding the health of women and their unborn children. ANC provide with preventive health care as well as pregnant women know from trained healthcare personnel regarding good healthy measures during the period from pregnancy, and better knowledge of danger signs during this period from pregnancy and childbirth, and receive social, sympathetic and psychological care at this vital time in their lives. The ANC services and institutional deliveries are significant to assess the condition of maternal care. Although home deliveries declined during this period of observation but at the same time the institutional deliveries also decreased which are matters of great concern.

The data analysis results regarding RCH services performance in the state of Rajasthan during the period from the month of April 2020 to June 2020 (during the period from lockdown) as compared to previous year 2019 (no lockdown) for same months suggest that the immunisation services had been adversely affected during the period from the lockdown period in 2020. It is evident from the findings of the study that not only newborn children but also the older ones have not received basic immunisation services.

Due to increasing population, there is a general trend of increase in need assessed as well as numbers achieved every year in all above indicators every year. The decline in achievement of such important indicators clearly signifies that SARS-CoV-2 pandemic have a negative effect on provision of RC health services such as maternal and child health. The role of family planning is very important in context of Rajasthan with increasing population and scanty natural resources. All family planning programmes reduced from previous year timeline for the same months indicating that the population control strategy of Rajasthan also suffered during this period.

<b>Immunisation</b>	<b>Item Code</b>	<b>Numbers of Achievement during the Period from April to June -2020-21</b>	<b>Numbers of Achievement during the Period from April to June -2019-20</b>	<b>Increase/Decrease in 2020 Compared to 2019 Percent</b>
BCG-Bacillus Calmette Guerin	9.1.2	263,111	307,840	Decrease 14.5%
PENTA 3 + DPT 3	9.1.5,9.1.8	215,897	319,423	Decrease 32.4%
DT (2 <sup>nd</sup> Dose) Or DPT-5	9.5.2	110,328	167,383	Decrease 34.1%
Measles + MR	9.2.1,9.2.2	313,047	350,584	Decrease 10.7%
Polio (OPV 3)	9.1.12	216,018	319,713	Decrease 32.4%
Tetanus Toxoid (TT10)	9.5.3	75,961	128,091	Decrease 40.7%
Tetanus Toxoid (TT16)	9.5.4	65,078	115,958	Decrease 43.9%
Vitamin A (1 <sup>st</sup> Dose)	9.8.1	306,922	340,687	Decrease 9.9%
Vitamin A (5 <sup>th</sup> )	9.8.2	122,037	318,123	Decrease 61.6%
Vitamin A (9 <sup>th</sup> )	9.8.3	66,533	293,822	Decrease 77.4%

**Table 4:** Comparison of immunisation coverage during 2019 and 2020 during the month of April to June (Rajasthan).

### ***Immunisation Coverage for 2019 and 2020 during the Month of April to June (Rajasthan)***

Table 4 shows that as compared to previous year 2019 data - For BCG immunisation there is a decline of 14.5 percent in 2020, Since BCG is given at birth it seems that either less number of children's is born during the period from this period or many have not received it, For Penta3+DPT3 the decline was 32.4 percent in 2020, For DT or DPT5 the decline was 34.1 percent in 2020, For Measles + MR the decline was 10.7 percent in 2020, For OPV3 the decline was 32.4 percent in 2020, For TT10 the decline was 40.7 percent in 2020, For TT16 the decline was 43.9 percent in 2020, For vitamin A which is given from 9 months of age 1<sup>st</sup> dose declined by 9.9 percent in 2020, Vitamin A 2<sup>nd</sup> to 9<sup>th</sup> dose starts from 16 months of age (one dose every 6 months), Vitamin A 5<sup>th</sup> dose declined by

61.6 percent in 2020, Vitamin A 9<sup>th</sup> dose declined by 77.4 percent in 2020. Here it is important to mention that vitamins A and OPV both are given orally. OPV3 is given at the age of 14 weeks whereas vitamin A is started from 9 month onwards. This observation suggests that the immunisation coverage for children of all ages declined in 2020 during April/May/June as compared to 2019 when the pandemic was not in existence.

Immunisation	Item Code	Need Assessed 2020-21	Achievement of Need Assessed 2020-21 during the Period from April to June	Percent Achievement of Need Assessed during the Period from April to June 2020-21	Need Assessed 2019-20	Achievement of-Need Assessed 2019-20 during the Period from April to June	Percent Achievement of Need Assessed during the Period from April to June 2019-20
BCG-Bacillus Calmette Guerin	9.1.2	1822630	263,111	14.4	1789690	307,840	17.2
PENTA 3+DPT 3	9.1.5,9.1.8	1822630	215,897	11.8	1789690	319,423	17.8
DT (2 <sup>nd</sup> Dose) OR DPT-5	9.5.2	1725000	110,328	6.4	1688500	167,383	9.9
Measles + MR	9.2.1,9.2.2	1822630	313,047	17.2	1789690	350,584	19.6
Polio (OPV 3)	9.1.12	1822630	216,018	11.9	1789690	319,713	17.9
Tetanus Toxoid (TT10)	9.5.3	1615140	75,961	4.7	1581000	128,091	8.1
Tetanus Toxoid (TT16)	9.5.4	1863310	65,078	3.5	1823900	115,958	6.4
Vitamin A (1 <sup>st</sup> Dose)	9.8.1	1822630	306,922	16.8	1789690	340,687	19.0
Vitamin A (5 <sup>th</sup> )	9.8.2	1780150	122,037	6.9	1742500	318,123	18.3
Vitamin A (9 <sup>th</sup> )	9.8.3	1725000	66,533	3.9	1688500	293,822	17.4


**Table 5:** Needs assessment and achievement comparison for immunisation.

#### ***Needs Assessment and Achievement Comparison for Immunisation***


Table 5 shows that as compared to previous year 2019 data - The need assessed for BCG (Bacillus Calmette Guerin) in 2020-2021 increased as compared to 2019-2020 (1822630-1789690) by 32940 numbers. The achievement of-need assessed in 2020-2021 during the period from April to June (263,111/14.4%) decreased as compared to 2019-2020 (307840/17.2%) by 44729 numbers and 2.8%; the need assessed for PENTA 3+DPT 3 in 2020-2021 increased as compared to 2019-2020 (1822630-1789690) by 32940 numbers. The achievement of-need assessed in 2020-2021 during the period from April to June (215897/11.8%) decreased as compared to 2019-2020 (319423/17.8%) by 103526 numbers and 6%; the need assessed for DT (2<sup>nd</sup> Dose) OR DPT-5 in 2020-2021 increased as compared to 2019-2020 (1725000-1688500) by 36500 numbers. The achievement of-need assessed in 2020-21 during the period from April to June (110328/06.4%) decreased as compared to 2019-2020 (167383/09.9%) by 57055 numbers and 3.5% (Figure 1 - Figure 3).

The need assessed for Measles + MR in 2020-2021 increased as compared to 2019-2020(1822630-1789690) by 32940 numbers. The achievement of-need assessed in 2020-2021 during the period from April to June (313047/17.2%) decreased as compared to 2019-20(350584/19.6%) by37537 numbers and 2.4%; the need assessed for POLIO (OPV 3) in 2020-2021 increased as compared to 2019-2020 (1822630-1789690) by 32940 numbers. The achievement of-need assessed in 2020-2021 during the period from April to June (216018/11.9%) decreased as compared to 2019-20(319713/17.9%) by 103695 numbers and 6%; the need assessed for Tetanus Toxoid (TT10) in 2020-2021 increased as compared to 2019-2020 (1615140-1581000) by 34140 numbers. The achievement of-need assessed in 2020-2021 during the period from April to June (75961/04.7%) decreased as compared to 2019-2020 (128091/08.1%) by52130 numbers and 3.4%; the need assessed for Tetanus Toxoid


(TT16) in 2020-2021 increased as compared to 2019-2020 (1863310-1823900) by 39410 numbers. The achievement of-need assessed in 2020-2021 during the period from April to June (65078/03.5%) decreased as compared to 2019-2020 (115958/06.4%) by 50880 numbers and 2.9%; the need assessed for Vitamin A (1<sup>st</sup> Dose) in 2020-2021 increased as compared to 2019-2020 (1822630-1789690) by 32940 numbers. The achievement of-need assessed in 2020-2021 during the period from April to June (306922/16.8%) decreased as compared to 2019-2020 (340687/19.0%) by 33765 numbers and 2.2%; the need assessed for Vitamin A (5<sup>th</sup> Dose) in 2020-2021 increased as compared to 2019-2020 (1780150-1742500) by 37650 numbers. The achievement of-need assessed in 2020-2021 during the period from April to June (122037/06.9%) decreased as compared to 2019-2020 (318123/18.3%) by 196086 numbers and 11.4%; the need assessed for Vitamin A (9<sup>th</sup> Dose) in 2020-2021 increased as compared to 2019-2020 (1725000-1688500) by 36500 numbers. The achievement of-need assessed in 2020-2021 during the period from April to June (66533/03.9%) decreased as compared to 2019-2020 (293,822/17.4%) by 227289 numbers and 13.5%.



**Figure 1:** Immunisation coverage chart for comparison of 2019-2020 and 2020-2021 for Rajasthan.



**Figure 2:** Chart for comparison on need assessed and achievement 2019-2020 and 2020-2021 for immunisation-Rajasthan.



**Figure 3:** Comparison on percent achievement 2019-2020 and 2020-2021 of need assessed during the period from April to June for immunisation 2020-2021 and 2019-2020.


Maternal Health	Item Code	Numbers of Achievement during the Period from April to June 2020-21	Need Assessed 2020-21	Percent of Need Achieved during the period from April to June 2020-21	Numbers of Achievement during the Period from April to June -2019-20	Need Assessed 2019-20	Percent of Need Achieved during the Period from April to June 2019-20	Percent Increase/Decrease in (Numbers of Achievement) 2020-21 Compared to 2019-20
ANC	1.1	445,804	2066780	21.6%	461,360	2031500	22.7%	Decrease 3.4%
Maternal Health Home Deliveries	2.1.1.A,2.1.1.B	6,489			5,588			Increase 16.1%
Maternal Health Institutional Deliveries	2.2	257,123	1892660	13.6%	299,624	1860380	16.1%	Decrease 14.2%

**Table 6:** Maternal health coverage during 2019-2020 and 2020-2021 during the month of April to June (Rajasthan).


**Maternal Health Coverage during 2019-20 and 2020-21 during the Month of April to June (Rajasthan)**

Regarding Maternal health coverage during 2019 and 2020 during the month of April to June (Rajasthan), Table 6 analyses shows that need assessed for ANC in 2020-2021 was 2066780 whereas the numbers of achievement during the period from April to June 2020-2021 was 445804 which was 21.6 percent of need assessed. In comparison need assessed for ANC in 2019-2020 was 2031500 whereas the numbers of achievement during the period from April to June 2019-2020 was 461,360 which was 22.7 percent of need assessed, the percent decrease in numbers of achievement during the period from 2020-2021 compared to 2019-2020 found a decrease of 3.4 percent in 2020-2021; The need assessed for home deliveries in 2020-2021 is not available whereas the numbers of achievement during the period from April to June 2020-2021 was 6489, In comparison need assessed for ANC in 2019-2020 is also not available whereas the numbers of achievement during the period from April to June 2019-2020 was 5588. The percent increase in numbers of achievement during the period from 2020-2021 compared to 2019-2020 found an increase of 16.1 percent in 2020-2021; The need assessed for institutional deliveries in 2020-2021 was 1892660 whereas the numbers of achievement during the period from April to June 2020-2021 was

257123 which was 13.6 percent of need assessed. In comparison need assessed in 2019-2020 were 1860380 whereas the numbers of achievement during the period from April to June 2019-2020 was 299624 which were 16.1 percent of need assessed. The percent decrease in numbers of achievement during the period from 2020-2021 compared to 2019-2020 found a decrease of 14.2 percent in 2020-2021 (Figure 4 and Figure 5).



**Figure 4:** Maternal health coverage for 2019-2020 and 2020-2021 during the month of April to June (Rajasthan).



**Figure 5:** Chart for comparison on need assessed and achievement 2019-2020 and 2020-2021 for maternal health-Rajasthan.

**Family Planning Coverage Comparison for 2019 and 2020 during the Month of April to June (Rajasthan)**

Regarding Family planning coverage comparison for 2019 and 2020 during the month of April to June (Rajasthan), Table 7 analyses shows that- the need assessed for condom user in 2020-2021 was 8995120 whereas need assessed for condom user in 2019-2020 was 8740800, an increase of 254320 is seen in 2020-2021; the numbers of achievement during the period from April to June 2020-2021 was 419235 which was 04.7 percent of need assessed whereas the numbers of achievement during the period from April to June 2019-2020 was 449495 which was 05.1 percent of need assessed. The Percent decrease in numbers of achievement during the period from April to June 2020-2021 compared to 2019-2020 found a decrease of 06.7 percent in 2020-2021. The need

assessed for oral pill users in 2020-2021 was 8995120 whereas need assessed for oral pill user in 2019-2020 was 8740800, an increase of 254320 is seen in 2020-2021; the numbers of achievement during the period from April to June 2020-2021 was 235,582 which was 02.6 percent of need assessed whereas the numbers of achievement during the period from April to June 2019-2020 was 252,298 which was 02.9 percent of need assessed. The percent decrease in numbers of achievement during the period from 2020-2021 compared to 2019-2020 found a decrease of 06.6 percent in 2020-2021. The need assessed for IUD insertion in 2020-2021 was 8995120 whereas in comparison need assessed for IUD insertion in 2019-2020 was 8740800, an increase of 254320 is seen in 2020-2021; the numbers of achievement during the period from April to June 2020-2021 was 89,539 which was 01.0 percent of need assessed whereas the numbers of achievement during the period from April to June 2019-2020 was 124,786 which were 01.4 percent of need assessed. The percent decrease in numbers of achievement during the period from April to June 2020-2021 compared to 2019-2020 found a decrease of 28.2 percent in 2020-2021.

Family Planning	Item Code	Numbers of Achievement April to June 2020-21	Need Assessed 2020-21	Percent of Need Achieved during the Period from April to June 2020-21	Numbers of Achievement April to June 2019-20	Need Assessed 2019-20	Percent of Need Achieved during the Period from April to June 2019-20	Percent Increase /Decrease in Numbers of Achievement 2020-21 Compared to 2019-20
Condom User	8.13	419,235	8995120	4.7	449,495	8740800	5.1	Decrease 6.7%
Oral Pill User	8.12, 8.14	235,582	8995120	2.6	252,298	8740800	2.9	Decrease 6.6%
IUD Insertion	8.3,8.4,8.5	89,539	8995120	1.0	124,786	8740800	1.4	Decrease 28.2%
Sterilisation	8.2.1,8.2.2, 8.2.3,8.2.4,8.1.1	13,698	15220170	0.1	61,399	14789800	0.4	Decrease 77.7%
Tubectomy	8.2.1,8.2.2, 8.2.3,8.2.4,	13,561	15220170	0.1	60,768	14789800	0.4	Decrease 77.7%
Vasectomy	8.1.1	137	15220170	0.0	631	14789800	0.0	Decrease 78.3%

**Note:** Sterilisation includes both tubectomy and vasectomy but need to be shown separately as per international guidelines and moreover the two processes are also different, tubectomy is done for female sterilisation whereas vasectomy is done for male sterilisation. The percentage of achievement of need assessed is different from the percentage comparison of numbers of achievement during the period from April to June of two financial years.

**Table 7:** Family planning coverage comparison for 2019 and 2020 during the period from the month of April to June (Rajasthan).

The need assessed for sterilisation (Figure 6 - Figure 9) in 2020-2021 was 15220170 whereas in comparison need assessed for sterilisation in 2019-2020 was 14789800, an increase of 430370 is seen in 2020-2021; the numbers of achievement during the period from April to June 2020-2021 was 13698 which was 0.1 percent of need assessed whereas the numbers of achievement during the period from April to June 2019-2020 was 61399 which were 0.4 percent of need assessed. The percent decrease in numbers of achievement during the period from April to June 2020-2021 compared to 2019-2020 found a decrease of 77.7 percent in 2020-2021. The need assessed for tubectomy in 2020-2021 was 15220170 whereas in comparison need assessed for tubectomy in 2019-2020 was 14789800, an increase of 430370 is seen in 2020-2021; the numbers of achievement during the period from April to June 2020-2021 was 13561 which was 0.1 percent of need assessed whereas the numbers of achievement during the period from April to June 2019-2020 was 60768 which were 0.4 percent of need assessed. The percent decrease in numbers of achievement during the period from 2020-2021 compared to 2019-2020 found a decrease of 77.7 percent in 2020-2021. The need assessed for vasectomy in 2020-2021 was 15220170 whereas in comparison need assessed for vasectomy in 2019-2020 was 14789800, an increase of 430370 is seen in 2020-2021; the numbers of

achievement during the period from April to June 2020-2021 was 137 which was 0.0 percent of need assessed whereas the numbers of achievement during the period from April to June 2019-2020 was 631 which were 0.0 percent of need assessed. The percent decrease in numbers of achievement during the period from 2020-2021 compared to 2019-2020 found a decrease of 78.3 percent in 2020-2021.




Figure 6: Chart for comparison on percent achievement 2019-2020 and 2020-2021 for maternal health- Rajasthan.




Figure 7: Family planning coverage comparison for 2019-2020 and 2020-2021, month of April to June -Rajasthan.





Figure 8: Comparison on need assessed for family planning coverage for 2019-2020 and 2020-2021-Rajasthan.



**Figure 9:** Comparison on percent achievement of need assessed for family planning coverage for 2019-2020 and 2020-2021.

## **DISCUSSION**

The observation, study, calculation and analysis of data obtained from HMIS (health management information system), Ministry of Health and Family Welfare website for RCH programme in Rajasthan shows that the lock down period and initial early phase of SARS-CoV-2 pandemic had a negative effect on the provision of RCH health services as detailed in results above. Besides the pandemic Rajasthan is also having lack of resources, skilled manpower poor infrastructure as well as positive deviance at community level [13]. These are the hindrances in the provision of RCH services beside the epidemic.

RCH (reproductive and child health) service provision is always a priority and great concern for populous nations particularly India with high levels of fertility. During the period from the SARS-CoV-2 pandemic era it's more challenging to deliver such essential services due to fear factors at community as well as health personal level. Fall in rate of immunisation can lead to emergence of diseases which is being controlled by immunisation programmes. If an unvaccinated child is open to the elements to disease microorganisms, the child's body may not be physically powerful as much as necessary to fight the disease. Before the discovery and invention of vaccines, several children died from diseases that vaccines now put off; such examples are whooping cough, measles, and polio. Though same microorganisms are present today, but since babies are cosseted by vaccines, we don't see these diseases nearly as frequently. Immunizing particularly children's also helps to guard the health of our community, in particular those who cannot be immunized (children who are too young to be vaccinated, or those who can't receive assured vaccines for medicinal/therapeutic reasons), and the little percentage of children who don't act in response to a particular vaccine. Vaccine-preventable diseases have an expensive brunt, resulting in hospital visits, hospitalizations, and early premature morbidity and mortality. Sick children can also cause parents to lose time from work [14].

Limited, insufficient and under quality ANC services will pose a threat to maternal as well as foetus life. Good quality antenatal care also provides pregnant women with micronutrient supplementation, management for hypertension to inhibit eclampsia, as well as immunization against tetanus. Antenatal care makes available HIV

testing and provides drugs to prevent mother-to-child transmission of HIV in pregnancy, childbirth, or breastfeeding (through breast milk). In malaria endemic zones, health personnel can make available pregnant women with drugs and insecticide-treated mosquito nets (ITNs) to help put off this devastating and sometimes deadly disease [15].

The insufficiency in family planning coverage will add more to burden of population explosion putting more pressure on available resources depletion. Women should be encouraged for making the choice, to take decision about the number and spacing of their conception. Family planning can help women protect their health and fertility and also in improving the quality of their own and family lives. Family planning also help in getting better children's health and ensuring access to enough food, clothing, housing, and educational opportunities. Family planning achieves these improvements in health and value of life in a better way compared with fund investments in most other health and social interventions. Committing human and financial assets for improving family planning services will get better health and well-being of women and children, and at the same time it will also prop up efforts to achieve a sustainable global population [16].

All states and union territories of India have enforced various different strategies, but it appears to be not totally effective to get good results on RC Health programme. The Government of Rajasthan should consider taking help of local intellectual people and communities as well as increasing skilled human resources working to safeguard maternal and child health in situations of distress-emergency such as SARS-CoV-2 pandemic. For this the barriers in provision of RCH services like a smaller number of skilled doctors, nurses, other resources should be reduced-rectified and promoters of RCH programme like health promotion activities creating awareness of benefits and addressing misbelieves and questions needs to be considered by policy-decision makers to achieve the maximum output from efforts and available resources.

Observations and analysis based on HMIS data from Bihar, Uttar Pradesh, Andhra Pradesh and India has also been conducted and submitted for peer reviewed publications which is available online for further reference [17-21].

The topic is of prime concern particularly for LMICs (low- and middle-income countries). Some researchers have used one modern LiST (lives saved tool) method to calculate the impact. One such study done by Timothy Robertson et al, found the indirect impacts of the SARS-COV-2 pandemic on maternal and child mortality in LMICs in which the health service coverage of essential maternal and child health interventions is reduced by 9.8%-51.9% and the prevalence of wasting is up by 10%-50% [22].

## **LIMITATIONS**

The data although collected from an accredited source may have some errors of reporting and human error etc. Moreover, it's stated by government that they use to refresh the data if some corrections are required. If the data is changed by the government which is usually rare the results will change accordingly.

## **CONCLUSION**

India and different states have launched several plans and strategies to deal with SARS-CoV-2 pandemic. However, this study report found that insufficient attention has been given to very essential services of RCH in

Rajasthan as evident from the results of this study. The problems in the provision of healthcare services should be dealt properly added with a proper update latest dynamic plan to carry on essential health delivery services even in pandemics and other situations of emergencies. The Rajasthan should develop an exclusive updated latest dynamic plan to deal with situations to guarantee provision of very essential services such as RCH during the period from pandemics or any other natural calamities-emergencies.

The ongoing SARS-CoV-2 pandemic has disclosed the vulnerability of women and children's and at the same time teaches us about importance of equity in healthcare. The children and women's who are considered most vulnerable in emergencies and distress must have special supportive healthcare all the time especially during the period from pandemics-emergencies and other natural disasters.

India is the second most populous country in the world after China. To control the ongoing population explosion including Rajasthan all states should have a robust population control strategy to working in all the conditions of emergency as well as in normal conditions.

### **COMPETING INTERESTS**

The authors declare that they have no competing interests in this section.

### **REFERENCES**

1. Karan A, Negandhi H, Hussain S et al. (2021) Size, composition and distribution of health workforce in India: Why, and where to invest?. *Human Resources for Health* 19(1): 1-14.
2. Kumar (2021) What impact have SARS-CoV-2/Covid-19 pandemic induced lockdown on the number of opd patients of diabetes, hypertension, stroke (cva), acute heart disease, mental illness, epilepsy, ophthalmic, dental and oncology in India during the lockdown months (April-May-2020) - observational research analysis?.
3. Reid A (2005) The effects of the 1918-1919 influenza pandemic on infant and child health in Derbyshire. *Medical History* 49(1): 29-54.
4. <http://www.nrhmhp.gov.in/content/immunisation>
5. <https://www.who.int/news/item/23-04-2021-covid-19-continues-to-disrupt-essential-health-services-in-90-of-countries>
6. <https://www.who.int/pmnch/covid-19/toolkits/child/child-protection/en>
7. [https://www.mohfw.gov.in/pdf/Annexure\\_MHA.pdf](https://www.mohfw.gov.in/pdf/Annexure_MHA.pdf)
8. Population census of India 2011.
9. (2020) World health organisation, Departmental news.
10. [http://rchiips.org/nfhs/factsheet\\_nfhs-4.shtml](http://rchiips.org/nfhs/factsheet_nfhs-4.shtml)
11. McClure EM, Kinney MV, Leisher SH et al. (2020) Impact of COVID-19 on maternal and child health. *The Lancet Global Health* 8(10): e1258.
12. Ministry of health and family welfare, Government of India.
13. Anand S, Fan V, World Health Organization (2016) The health workforce in India. World Health Organization.

14. <https://www.cdc.gov/vaccines/vac-gen/default.htm>
15. (2021) Antenatal care. UNICEF.
16. World Health Organization (1994) Mother-baby package: A safe motherhood planning guide, WHO, Geneva.
17. Piyush Kumar (2020) What impact have SARS-CoV-2/Covid-19 pandemic on the reproductive and child health programme of India over the 3 months after nationwide lock down announcement in March 2020 - A brief analysis.
18. Piyush Kumar (2021) What impact have SARS-CoV-2/Covid-19 pandemic on the reproductive and child health programme of Bihar in India over the 3 months after nationwide Lock down announcement in March 2020? How SARS-CoV-2 Pandemic era does influence RCH Programme, Immunisation, and Maternal Health?
19. Piyush Kumar (2021) What impact have SARS-CoV-2/Covid-19 pandemic on the reproductive and child health programme of Uttar Pradesh in India over the 3 months after nationwide lockdown announcement in March 2020 -A brief analysis.
20. Kumar P (2021) What impact have SARS-CoV-2/Covid-19 pandemic on the reproductive and child health programme (immunization, maternal health, family planning) of Andhra Pradesh in India over the 3 months after nationwide lock down announcement in March 2020?
21. Piyush Kumar (2021) What impact have SARS-CoV-2/Covid-19 pandemic on the reproductive and child health programme of India over the 3 months after nationwide lockdown announcement in March 2020 -A brief analysis.
22. Roberton T, Carter ED, Chou VB et al. (2020) Early estimates of the indirect effects of the COVID-19 pandemic on maternal and child mortality in low-income and middle-income countries: A modelling study. *The Lancet Global Health* 8(7): e901-e908.